

The Testate Lobose Amoebae (Order Arcellinida Kent, 1880) Finally Find their Home within Amoebozoa

Sergey I. Nikolaev^{a,d,1}, Edward A. D. Mitchell^{b,c}, Nikolay B. Petrov^a, Cédric Berney^d, José Fahrni^d, and Jan Pawlowski^d

^aDepartment of Evolutionary Biochemistry, A. N. Belozersky Institute of Physico-Chemical Biology, Moscow State University, Moscow, Russia

^bEcole Polytechnique Fédérale de Lausanne (EPFL), Faculté Environnement Naturel, Architectural et Construit, Laboratoire des Systèmes Écologiques, station 2, CH-1015 Lausanne, Switzerland

^cSwiss Federal Research Institute WSL, Antenne Romande, station 2, CH-1015 Lausanne, Switzerland

^dDepartment of Zoology and Animal Biology, University of Geneva, Sciences III 30, quai Ernest Ansermet CH-1211 Genève 4, Switzerland

Submitted December 20, 2004; Accepted March 1, 2005
Monitoring Editor: Michael Melkonian

Testate lobose amoebae (order Arcellinida Kent, 1880) are common in all aquatic and terrestrial habitats, yet they are one of the last higher taxa of unicellular eukaryotes that has not found its place in the tree of life. The morphological approach did not allow to ascertain the evolutionary origin of the group or to prove its monophyly. To solve these challenging problems, we analyzed partial small-subunit ribosomal RNA (SSU rRNA) genes of seven testate lobose amoebae from two out of the three suborders and seven out of the 13 families belonging to the Arcellinida. Our data support the monophyly of the order and clearly establish its position among Amoebozoa, as a sister-group to the clade comprising families Amoebidae and Hartmannellidae. Complete SSU rRNA gene sequences from two species and a partial actin sequence from one species confirm this position. Our phylogenetic analyses including representatives of all sequenced lineages of lobose amoebae suggest that a rigid test appeared only once during the evolution of the Amoebozoa, and allow reinterpretation of some morphological characters used in the systematics of Arcellinida.

© 2005 Published by Elsevier GmbH.

Key words: testate amoebae; phylogeny; evolution; SSU rRNA gene; actin; Amoebozoa.

Introduction

The testate lobose amoebae (order Arcellinida Kent, 1880) are common in a wide range of moist

and freshwater habitats. These small amoebae feed mostly on bacteria, algae, and fungi. The larger species also prey on other protozoans and small metazoans (Heal 1963; Mast and Root 1916). The Arcellinida are distinguished by their tests, comprising a single aperture and composed of

¹Corresponding author;
fax +41 22 379 67 95
e-mail s-nikol@yandex.ru (S.I. Nikolaev)

either secreted proteinaceous material or agglutinated. Proteinaceous tests can be flexible, with rigid sheets of fibrous material or with regularly arranged hollow building units that form an areolate surface. Agglutinated tests can be either calcareous or siliceous (Ogden 1990; Ogden and Ellison 1988). Dictyosomes are involved in the secretion of the organic building units and the cement. Mitochondria have branched tubular cristae. Contractile vacuoles are present. Arcellinida have either endolobopodia that are granular or completely hyaline, or ectolobopodia that are generally fingerlike and in some species can anastomose (reticulolobopodia) (Bonnet 1961, 1963). Under unfavorable environmental conditions, most testate lobose amoebae produce cysts.

Classification of the Arcellinida is based mainly on characters of the test. Kudo (1954) presents the arcellinids as two unrelated families, Arcellinidae (with a membranous shell) and Diffugiidae (shell with foreign bodies, platelets, or scales). A more recent classification (Meisterfeld 2002) considers Arcellinida as a monophyletic order divided into three suborders: Arcellinina (membranous test, digitate pseudopodia), Diffugiina (test rigid with mineral particles, digitate pseudopodia), and Phryganellina (test with siliceous material, pseudopodia conical).

Testate lobose amoebae were for a long time considered as part of the Testacea (Rhizopoda), a taxon uniting all amoeboid protists that are enveloped by a single-chambered shell (Kudo 1954). In the first edition of the "Illustrated Guide to Protozoa" (Bovee 1985), the Testacea are not treated as a monophyletic taxon. The shape of pseudopodia (lobose or filose) is considered as an important taxonomic feature and testate lobose amoebae are placed within the class Lobosea Carpenter, 1861; while the testate filose amoebae are placed within the class Filosea Leidy, 1879. In the second edition of the "Illustrated Guide to the Protozoa" (Meisterfeld 2002), the Arcellinida are treated as a group of amoebae of uncertain affinities, because their evolutionary origins are unclear and molecular data are awaited to solve the problem.

Molecular data have already helped to resolve the phylogenetic status of testate filose amoebae which form at least three independent lineages within the recently defined super-group Rhizaria (Burki et al. 2002; Nikolaev et al. 2003, 2004; Wylezich et al. 2002). In contrast, the Arcellinida is one of the last widespread and well-known groups of eukaryotes that is not represented in molecular databases. To test the monophyly of Arcellinida,

determine their position in the eukaryotic tree, and to resolve their internal relationships, we obtained partial small-subunit ribosomal RNA (SSU rRNA) gene sequences from representatives of two out of the three suborders and seven out of the 13 families of Arcellinida. We show that testate lobose amoebae are monophyletic and belong to the Amoebozoa, and confirm this result using complete SSU rRNA gene sequences from two species, and a partial actin sequence from one species.

Results

Seven species of testate lobose amoebae (*Arcella artocrea*, *Bullinularia indica*, *Centropyxis laevigata*, *Heleopera sphagni*, *Hyalosphenia papilio*, *Nebela tinctoria* var. *major*, and *Trigonopyxis arcuata*) were examined (Fig. 1, Table 1). The partial SSU rRNA gene tree (Fig. 2) shows that the seven species form a monophyletic group within Amoebozoa. Monophyly of the Arcellinida is supported with both MrBayes and Maximum Likelihood analyses. The relationships within Arcellinida are relatively well resolved. *Heleopera sphagni* branches at the base of the testate lobose amoebae with strong support (PP of 1.00 and BV of 82%). The next branch is comprised of *H. papilio* and *N. tinctoria* var. *major*, which group together with high support (PP of 1.00 and BV of 100%). The four remaining species group together with moderate support (PP of 0.82 and BV of 64%) and are divided in two highly supported (PP of 1.00 and BV of 100%) clusters: *T. arcuata*+*B. indica* and *A. artocrea*+*C. laevigata*.

Analysis of complete SSU rRNA sequences (Fig. 3), including two arcellinid species (*H. sphagni* and *C. laevigata*), further supports the monophyly of Arcellinida, recovered here with strong support (PP of 0.97 and BV of 83%). In both SSU rRNA gene trees (Figs 2, 3), the testate lobose amoebae branch within a highly supported clade (PP of 1.00, BV of 94–100%) comprising the genus *Echinamoeba*, the order Leptomyxida, and the families Hartmannellidae and Amoebidae. This clade was named Tubulina in a recent classification of lobose amoebae (Smirnov et al. 2005), and we follow this classification here. Within Tubulina, a close relationship was recovered between testate lobose amoebae, and the families Amoebidae and Hartmannellidae. Support for this relation is moderate in the partial SSU rRNA analysis (PP of 0.63 and BV of 68%; Fig. 2), but it is high using complete SSU rRNA sequences




Figure 1. Micrographs of the seven testate amoebae taxa analyzed: **a, b:** *Heleopera sphagni* (small spheres inside the cyst are endosymbiotic algae); **c:** *Nebela tinctoria* var. *major*; **d, e:** *Hyalosphenia papilio*; **f:** *Bullinularia indica*; **g:** *Trigonopyxis arcuata*; **h:** *Arcella artocrea*; **i:** *Centropyxis laevigata*. Scale bars indicate 50 μm . SEM pictures (b, d, g, and i) were taken at the University of Alaska Anchorage by Drs. J. Kudenov, E. Mitchell, and K. Kishaba.

(PP of 1.00, BV of 92%; Fig. 3). All other well-supported amoebozoan clades present in the SSU rRNA trees are congruent with previous studies and will not be discussed here.

The actin phylogeny (Fig. 4) corroborates the results obtained with the SSU rRNA data. A strong relationship (PP of 1.00 and BV of 100%) is recovered between Arcellinida and members of the families Amoebidae and Hartmannellidae. However, the resolution between the different clades of lobose amoebae is poor, and the lack of actin sequences for the basal members of the Tubulina impedes more precise conclusions about the position of Arcellinida within this clade.

Discussion

Up to now, the Arcellinida remained one of the last higher taxa of eukaryotes for which no molecular data were available. The classical morphological approach failed to ascertain the evolutionary origin of this group so far (Meisterfeld 2002). The idea of placing testate lobose amoebae (order Arcellinida) within the group of lobose amoebae, class Lobosea Carpenter, 1861 was put forward by Deflandre (1953) and Loeblich and Tappan (1961, 1964), who united all amoebae with lobose pseudopodia. Results of our analyses clearly confirm that testate lobose amoebae are indeed

Table 1. Morphological characteristics of the seven Arcellinida taxa studied.

	Shell dimensions					Shell composition ^a	Position of aperture ^b	References for dimensions	Sampling location
	Diameter [μm]	Length [μm]	Breadth [μm]	Depth [μm]	Diameter of aperture [μm]				
<i>Arcella artocrea</i> Deflandre, 1928	184–216			46–64	36–42	S	V	Ogden and Hedley 1980	Creux de l'Epral (CH)
<i>Bullinularia indica</i> Penard, 1907	138–180		165–172	94–99	65–90	A	V	Ogden and Hedley 1980	Creux de l'Epral (CH)
<i>Centropyxis laevigata</i> Penard, 1902	70–135					A	V	Deflandre 1929	Anchorage (AK, USA)
<i>Heleopera sphagni</i> Leidy, 1874		80–145	50–120	42–51	40–45	A	T	Ogden and Hedley 1980	Anchorage (AK, USA)
<i>Hyalosphenia papilio</i> Leidy, 1875		90–175	60–115	21–25	32–40	S	T	Ogden and Hedley 1980	Anchorage (AK, USA)
<i>Nebela tinctoria</i> var. <i>major</i> Deflandre, 1936		90–120				R	T	Deflandre 1936	Creux de l'Epral (CH)
<i>Trigonopyxis arcuata</i> Leidy, 1879	90–168			40–80	21–45	A	V	Ogden and Hedley 1980	Anchorage (AK, USA)

^aS: secreted proteinaceous; A: agglutinate; R: recycled biosilica,

^bT: terminal aperture; V: ventral aperture




Figure 2. MrBayes tree of partial SSU rRNA genes showing the monophyly of Arcellinida. Numbers at nodes indicate posterior probabilities for MrBayes analysis (upper) and bootstrap values for ML analysis (down). The scale bar indicates 0.1% sequence divergence.




Figure 3. MrBayes tree of full length SSU rRNA genes indicating the phylogenetic position of Arcellinida among Amoebozoa. Numbers at nodes indicate posterior probabilities for MrBayes analysis (upper) and bootstrap values for ML analysis (down). The scale bar indicates 0.1% sequence divergence.




Figure 4. MrBayes tree of actin genes indicating the phylogenetic position of Arcellinida among Amoebozoa. Numbers at nodes indicate posterior probabilities for MrBayes analysis (upper) and bootstrap values for ML analysis (down). The scale bar indicates 0.02% sequence divergence.

related to naked lobose amoebae. The Arcellinida belong to the Amoebozoa, a position supported by both SSU rRNA and actin gene data (Figs 2–4). Members of the Arcellinida possess branching tubular mitochondrial cristae (Meisterfeld 2002), as do other lobose amoebae, which were named ramicristate amoebae by Rogerson and Patterson (2002). This suggests that the shape of mitochondrial cristae might be a valid derived feature of the Amoebozoa.

Both SSU rRNA and actin gene data support a close relationship between testate lobose amoebae and the families Amoebidae and Hartmannellidae. This is in agreement with the hypothesis, based on the shape of lobopodia, that the Arcellinida are related to typical large lobose amoebae such as *Amoeba proteus*, with poly-podial or monopodial locomotion, and possessing classical fingerlike, non-eruptive pseudopodia, cylindrical in cross-section (Smirnov et al. pers. commun.).

Based on morphological characters alone, it was supposed that Arcellinidae species with a

simple membranous shell have a different origin from the species with foreign bodies and scales incorporated in the shell (Kudo 1954). Our analyses of partial SSU rRNA sequences including seven species of testate lobose amoebae belonging to two out of the three suborders and seven out of the 13 families of Arcellinida indicate that the order Arcellinida is monophyletic (Fig. 2). However, our results show that the character of the composition of the shell does not reflect the relationships within Arcellinida: in our analysis, species with proteinaceous (*Hyalosphenia* and *Arcella*) and agglutinated (*Nebela*, *Trigonopyxis*, *Centropyxis*, *Bullinularia*, and *Heleopera*) shells are mixed (Fig. 2). Our molecular phylogeny of the Arcellinida suggests that primitive members of the order probably started building shells by agglutinating organic and/or mineral debris from the environment (xenosomes), while some more evolved taxa (e.g. *Nebela*) later started producing shells by recycling biosilica plates synthesized by their prey, such as filose testate amoebae from the order Euglyphida, or

produced proteinaceous shells (e.g. *Hyalosphenia* and *Arcella*). The independent branching of *Hyalosphenia* and *Arcella* provides evidence that the evolution towards more resistant proteinaceous shells took place more than once within the Arcellinida.

Given the topology shown in Figure 2, other morphological characters seem to make sense regarding the internal relationships of testate lobose amoebae. The phylogenetic position of the seven studied taxa suggests that all taxa with a ventral aperture (*Arcella*, *Centropyxis*, *Bullinularia* and *Trigonopyxis*) share a common ancestry regardless of the shell composition (PP of 0.82 and BV of 64%, Fig. 1). In contrast, the taxa that produce shells with a terminal aperture form two successive lineages (*Heleopera* and *Hyalosphenia+Nebela*) at the base of the Arcellinida. The latter lineage shares very similar shell morphologies (Fig. 1), despite the different material used for shell construction. Based on these morphological similarities, these two genera were grouped in the family Hyalospheniidae, and this grouping is strongly supported by our results (PP of 1.00, BV of 100%). In contrast, *Heleopera* species use xenosomes to build their shells and in this respect they resemble some of the taxa with ventral aperture included in this study (*Centropyxis*, *Bullinularia*, and *Trigonopyxis*).

Our data provide some support to a hypothetical phylogeny based on morphology (Bonnet 1964), suggesting that a test with a terminal aperture is a primitive trait within the Arcellinida, while a test with a ventral aperture is a derived trait. Although more data are clearly needed to confirm this observation, it also would make sense in light of the probable aquatic origin of the Arcellinida. Indeed, a ventral aperture, especially when combined with a flat shell, allows the amoeba to remain active in relatively dry conditions as long as a capillary water film is present on the substrate. All other things being equal, taxa with a ventral aperture can reach a larger volume and biomass under hydric stress than taxa with terminal apertures. In support of this idea, large *Nebela* species are only found in the wettest micro-sites in *Sphagnum*-dominated peatlands, while small taxa such as *N. militaris* and *N. tincta* are common in dry habitats such as hummocks and non-*Sphagnum* mosses, in places where much larger taxa with a ventral aperture such as *Trigonopyxis arcula* and *Bullinularia indica* occur regularly (Booth 2002; Charman and Warner 1992; Lamentowicz and Mitchell in press; Mitchell et al. 1999).

Methods

Testate amoebae were obtained from *Sphagnum* samples collected at two *Sphagnum*-dominated peatlands: Bicentennial park, Anchorage, Alaska (samples taken in fall 2003; 61°06'N, 149°44'W) and Le Creux de l'Epral, in the Swiss Jura Mountains (samples taken on September 15th 2004; 47°12'N, 4°37'E) (Table 1, Fig. 1). The specimens were extracted from the mosses and picked using a narrow diameter pipette under the dissecting microscope. Identifications were confirmed under the microscope at 400× magnification.

DNA was extracted using the DNeasy Plant Minikit (Qiagen, Basel, Switzerland) or with guanidine buffer (Chomczynski and Sacchi 1987). Partial SSU rRNA genes were amplified for *Arcella artocrea* (550 bp), *Hyalosphenia papilio* (999 bp), *Nebela tincta* var. *major* (670 bp), *Trigonopyxis arcula* (1762 bp), and *Bullinularia indica* (1076 bp), using the universal primers s12.2 (5'-GAT(CT)AGATACCGTCGTAGTC-3') and sB (5'-TGATCCTTCTGCAGGTTACCTAC-3'). Complete SSU rRNA gene sequences were obtained for *Heleopera sphagni* (2379 bp) and *Centropyxis laevigata* (2133 bp). The partial sequences amplified with s12.2 and sB were extended in the 5' direction using reverse primers specifically designed for each species paired with the forward universal primers sA (5'-ACCTGGTTGATCCTGC-CAGT-3') and s6 (5'-CNGCGGTAATTCCAGCTC-3'). A partial fragment (1093 bp) of the actin gene of *H. sphagni* was amplified using the forward primers Act-F1 (5'-CNGARGCDDCATTRAAYC-3') and Act-N2 (5'-AACTGGGA(CT)GA(CT)ATGGA-3') and the reverse primer Act-1354r (5'-GGACCA-GATTCATCATAYTC-3'). The fragment contains an intron of 311 bp and the amplified coding region is 782 bp long. PCR amplifications were carried out in a total volume of 50 µl with an amplification profile consisting of 40 cycles with 30 s at 94 °C, 30 s at 50 °C, and 2 min at 72 °C, followed by 5 min at 72 °C for the final extension. The amplified PCR products were purified using the High Pure PCR Purification Kit (Roche, Rotkreuz, Switzerland), then ligated into pGEM-T Vector System (Promega, Wallisellen, Switzerland), cloned in XL-2 Ultracompetent Cells (Stratagene, Basel, Switzerland), sequenced with the ABI-PRISM Big Dye Terminator Cycle Sequencing Kit, and analyzed with an ABI-3100 DNA sequencer (Perkin-Elmer, Rotkreuz, Switzerland), all according to the manufacturers' instructions. The sequences obtained in this study have been submitted to GenBank

Table 2. Taxonomic position, species names and GenBank accession numbers of all SSU rRNA and actin gene sequences used in this study.

Taxonomic position	Species name	GenBank accession numbers
New sequences presented in this study		
SSU rRNA gene		
Arcellinida	<i>Heleopera sphagni</i>	AY848964
	<i>Nebela tinctoria</i> var. <i>major</i>	AY848968
	<i>Hyalosphenia papilio</i>	AY848966
	<i>Bullinularia indica</i>	AY848970
	<i>Trigonopyxis arcuata</i>	AY848967
	<i>Centropyxis laevigata</i>	AY848965
	<i>Arcella artocrea</i>	AY848969
Actin gene		
Arcellinida	<i>Heleopera sphagni</i>	AY848971
Other sequences used in our analyses		
SSU rRNA gene		
Tubulina	<i>Echinamoeba exundans</i>	AF293895
	<i>Echinamoeba thermanum</i>	AJ489268
	<i>Hartmannella vermiformis</i>	M95168
	Uncultured eukaryote clone BOLA868	AF372795
	<i>Leptomyxa reticulata</i>	AF293898
	<i>Paraflabellula reniformis</i>	AF293900
	<i>Rhizamoeba saxonica</i>	AY121847
	<i>Paraflabellula hoguae</i>	AF293899
	Hartmannellidae sp. LOS7N/I	AY145442
	<i>Saccamoeba limax</i>	AF293902
	<i>Glaeseria mira</i>	AY294196
	<i>Hartmannella cantabrigiensis</i>	AY294147
	<i>Amoeba leningradensis</i>	AJ314605
	<i>Chaos nobile</i>	AJ314606
	<i>Amoeba proteus</i>	AJ314604
	<i>Chaos carolinense</i>	AJ314607
Vannellina+Conopodina	<i>Lingulamoeba leei</i>	AY183886
	<i>Vannella anglica</i>	AF099101
	<i>Platyamoeba placida</i>	AY294150
	<i>Platyamoeba plurinucleolus</i>	AY121849
	<i>Vexillifera armata</i>	AY183891
	<i>Neoparamoeba aestuarina</i>	AY121848
	<i>Korotnevella hemistylolopsis</i>	AY121850
	<i>Korotnevella stella</i>	AY183893
Acanthopodina	<i>Balamuthia mandrillaris</i>	AF019071
	<i>Acanthamoeba tubiashi</i>	AF019065
Other Amoebozoa	<i>Dermamoeba algensis</i>	AY294148
	<i>Mayorella</i> sp.	AY294143
	LKM74	AJ130863
	<i>Platyamoeba stenopodia</i>	AY294144
	<i>Phalansterium solitarium</i>	AF280078
	<i>Gephyramoeba</i> sp.	AF293897
	Uncultured eukaryote clone RT5iin44	AY082989
	<i>Filamoeba nolandi</i>	AF293896
Opisthokonta	<i>Trichoplax adhaerens</i>	L10828
	<i>Schizosaccharomyces pombe</i>	X58056

Table 2. (continued)

Taxonomic position	Species name	GenBank accession numbers
Other sequences used in our analyses		
Actin gene		
Tubulina	<i>Chaos carolinense</i>	AY294157
	<i>Amoeba proteus</i>	AJ314604
	<i>Hartmannella cantabrigiensis</i>	AY294158
	<i>Glaeseria mira</i>	AY294155
Vannellina	<i>Platyamoeba placida</i>	AY294153
	<i>Vannella ebro</i>	AY294151
Acanthopodina	<i>Acanthamoeba castellanii</i>	V00002
Other Amoebozoa	<i>Thecamoeba similis</i>	AY294154
	<i>Mayorella</i> sp.	AY294152
	<i>Dermamoeba algensis</i>	AY294159
Opisthokonta	<i>Saccharomyces cerevisiae</i>	L00026
	<i>Schizosaccharomyces pombe</i>	Y00447

(see Table 2 for the species names and accession numbers of all sequences used in this study).

The partial SSU rRNA gene sequences from *H. sphagni*, *N. tinctoria* var. *major*, *H. papilio*, *B. indica*, *T. arcula*, *C. laevigata*, and *A. artocrea* were manually fitted to an general alignment of eukaryotic SSU rRNA gene sequences, based on a universal model of eukaryotic SSU rRNA secondary structure (Van de Peer et al. 2000). Preliminary analyses indicated that all sequences of Arcellinida belong to the phylum Amoebozoa (data not shown). Our alignment was thus reduced to 32 sequences for the phylogenetic analyses, including the 7 sequences of testate lobose amoebae and 25 sequences from other Amoebozoa (all fast-evolving members of Amoebozoa that are clearly not closely related to testate lobose amoebae according to the preliminary analyses, such as archamoebae and mycetozoans, were omitted to allow the use of a maximum amount of nucleotide positions that was increased up to 606 bp). A second alignment of complete SSU rRNA gene sequences was prepared, including the sequences of *H. sphagni* and *C. laevigata*, 28 sequences from other Amoebozoa, and 2 sequences from opisthokonts used as outgroup. In the analyses 1623 positions were included. Finally, the partial actin sequence of *H. sphagni* was manually fitted to an alignment of 13 eukaryotic actin gene sequences, including 10 sequences of lobose amoebae and 2 sequences of Fungi used as outgroup. In the analyses 255 amino acid positions were included. Actin and SSU rRNA alignments are available upon request from the authors.

Maximum likelihood (ML) analyses were performed with PhyML, version 2.4 (Guindon and Gascuel 2003), using the GTR model of evolution (Lanave et al. 1984; Rodriguez et al. 1990) for all SSU rRNA gene analyses and the JTT model of evolution (Jones et al. 1992) for the amino acid alignment of actin gene sequences, taking into account a proportion of invariables sites, and a gamma distribution of the rates of substitution for the variable positions, with 4 rate categories. As starting trees, BIONJ trees were used. Bayesian analyses (BA) were performed with MrBayes (Huelsenbeck and Ronquist 2001) using the same models of evolution. The program was run for 1,000,000 generations, sampled every 100 generations, with 4 simultaneous chains. The trees sampled before the chains reached stationarity were discarded as a burn-in.

Acknowledgements

The authors thank Alexey Smirnov for the discussion of the work, and Ralf Meisterfeld and an anonymous reviewer for important corrections that improved the quality of the manuscript. This work was supported by the Russian Foundation for Basic Research 02-04-48265, 02-04-48958, 02-04-49987, 00-15-97905, and the Swiss National Science Foundation grants 3100-064073.00 and 7SUPJ062343. This work was initiated while Edward Mitchell was at the University of Alaska Anchorage and was completed while he was at EPFL and WSL-AR, and was supported by EU project RECIPE. RECIPE is partly supported by

the European Commission (n EVK2-2002-00269) and partly, for the Swiss partners EPFL and WSL-AR, by the OFES (Swiss Federal Office for Education and Science), Switzerland. The SEM pictures used in Fig. 1 were done at the University of Alaska Anchorage SEM Lab by Dr. Jerry Kudenov, Keiko Kishaba and Edward Mitchell.

References

- Bonnet L** (1961) L'émission pseudopodique chez les thécamoebiens endogés (I). *Bull Soc Zool Fr* **86**: 17–28
- Bonnet L** (1963) L'émission pseudopodique chez les thécamoebiens endogés (II). *Bull Soc Zool Fr* **88**: 57–63
- Bonnet L** (1964) Le peuplement thécamoebiens des sols. *Rev Écol Biol Sol* **1**: 123–408
- Booth RK** (2002) Testate amoebae as paleoindicators of surface-moisture changes on Michigan peatlands: modern ecology and hydrological calibration. *J Paleolimnol* **28**: 329–348
- Bovee EC** (1985) Order Foraminiferida D'Orbigny, 1826. In Lee JJ, Hutner SH, Bovee EC (eds) *An Illustrated Guide to the Protozoa*. Society of Protozoologists, Lawrence, Kansas, pp 158–211
- Burki F, Berney C, Pawlowski J** (2002) Phylogenetic position of *Gromia oviformis* Dujardin inferred from nuclear-encoded small subunit ribosomal DNA. *Protist* **153**: 251–260
- Charman DJ, Warner BG** (1992) Relationship between testate amoebae (Protozoa, Rhizopoda) and microenvironmental parameters on a forested peatland in Northeastern Ontario. *Can J Zool-Rev Can Zool* **70**: 2474–2482
- Chomczynski P, Sacchi N** (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction. *Anal Biochem* **162**: 156–159
- Deflandre G** (1929) Le genre *Centropyxis* Stein. *Arch Protistenkd* **67**: 322–375
- Deflandre G** (1936) Étude monographique sur le genre *Nebela* Leidy. *Ann Protistol* **5**: 201–286
- Deflandre G** (1953) Ordres des Testaceolobosa (de Saedeleer, 1934), Testaceafilosa (de Saedeleer, 1934), Thalamia (Haeckel, 1862) ou Thécamoebiens (Auct.) (Rhizopoda Testacea). In Grassé P-P (ed) *Traité de Zoologie*, vol. 1. Masson and Co., Paris, pp 97–148
- Guindon S, Gascuel O** (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* **52**: 696–704
- Heal OW** (1963) The distribution of testate amoebae (Rhizopoda, testacea) in some fens and bogs in Northern England. *J Linn Soc (Zool)* **23**: 254–264
- Huelsenbeck JP, Ronquist F** (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**: 754–755
- Jones DT, Taylor WR, Thornton JM** (1992) The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci* **8**: 275–282
- Kudo RR** (1954) *Protozoology*. Charles C. Thomas, Springfield, IL, pp 472–492
- Lamentowicz M, Mitchell EAD** in press. The ecology of testate amoebae (Protists) in *Sphagnum* in north-west Poland in relation to peatland ecology. *Microb Ecol*
- Lanave C, Preparata G, Saccone C, Serio G** (1984) A new method for calculating evolutionary substitution rates. *J Mol Evol* **20**: 86–93
- Loeblich AR, Tappan H** (1961) Suprageneric classification of the Rhizopoda. *J Paleontol* **35**: 245–330
- Loeblich AR, Tappan H** (1964) Foraminiferal facts, fallacies and frontiers. *Geol Soc Am Bull* **75**: 367–392
- Mast SO, Root FM** (1916) Observations on amoeba feeding on rotifers, nematodes and ciliates and their bearing on the surface-tension theory. *J Exp Zool* **21**: 33–49
- Meisterfeld R** (2002) Order Arcellinida Kent, 1880. In: Lee JJ, Leedale GF, Bradbury P (eds), *The Illustrated Guide to the Protozoa*. 2 ed., vol. 2, Allen Press, Lawrence, KS, pp 827–860
- Mitchell EAD, Buttler AJ, Warner BG, Gobat J-M** (1999) Ecology of testate amoebae (Protozoa: Rhizopoda) in *Sphagnum* peatlands in the Jura mountains, Switzerland and France. *Ecoscience* **6**: 565–576
- Nikolaev SI, Berney C, Fahrni JF, Mylnikov AP, Aleshin VV, Petrov NB, Pawlowski J** (2003) *Gymnophrys cometa* and *Lecythium* sp. are core Cercozoa: Evolutionary implications. *Acta Protozool* **42**: 183–190
- Nikolaev SI, Berney C, Fahrni JF, Bolivar I, Polet S, Mylnikov AP, Aleshin VV, Petrov NB, Pawlowski J** (2004) The twilight of Heliozoa and rise of Rhizaria, an emerging supergroup of amoeboid eukaryotes. *Proc Natl Acad Sci U S A* **101**: 8066–8071

Ogden CG (1990) The Structure of the Shell Wall in Testate Amoebae and the Importance of the Organic Cement Matrix. In Claugher D (ed) *Scanning Electron Microscopy in Taxonomy and Functional Morphology*. Systemat Assoc Special vol **41**. Clarendon Press, Oxford, pp 235–257.

Ogden CG, Ellison RL (1988) The value of the organic cement matrix in the identification of the shells of fossil testate amoeba. *J Micropaleontol* **7**: 233–240

Ogden CG, Hedley RH (1980) *An Atlas to Freshwater Testate Amoebae*. Oxford University Press, Oxford

Rogerson A, Patterson DJ (2002) Ramicristate Amoebae (Gymnamoebae). In Lee JJ, Leedale GF, Bradbury P (eds) *The Illustrated Guide to the Protozoa*. 2nd ed. Society of Protozoologists, Lawrence, KS, pp 1023–1052

Rodriguez F, Oliver JF, Martin A, Medina JR (1990) The general stochastic model of nucleotide substitution. *J Theor Biol* **142**: 485–501

Smirnov A, Nasonova E, Berney C, Fahrni J, Bolivar I, Pawlowski J (2005) Molecular phylogeny and classification of the Lobose amoebae. *Protist* **156**: 129–142

Van de Peer Y, De Rijk P, Wuyts J, Winkelmanns T, De Wachter R (2000) The European small subunit ribosomal RNA database. *Nucleic Acids Res* **28**: 175–176

Wylezich C, Meisterfeld R, Meisterfeld S, Schlegel M (2002) Phylogenetic analyses of small subunit ribosomal RNA coding regions reveal a monophyletic lineage of euglyphid testate amoebae (order Euglyphida). *J Eukaryot Microbiol* **49**: 108–118

Available online at www.sciencedirect.com

