

Cross-points in the Dirichlet-Neumann method I : well-posedness and convergence issues

Bastien Chaudet-Dumas^{1*} and Martin J. Gander¹

^{1*}Section de Mathématiques, Université de Genève, Geneva,
CH-1205, Switzerland.

*Corresponding author(s). E-mail(s): bastien.chaudet@unige.ch;
Contributing authors: martin.gander@unige.ch;

Abstract

Cross-points in domain decomposition, i.e. points where more than two subdomains meet, have received substantial attention over the past years, since domain decomposition methods often need special attention in their definition at cross-points, in particular if the transmission conditions of the domain decomposition method contain derivatives, like in the Dirichlet-Neumann method. We study here for the first time the convergence of the Dirichlet-Neumann method at the continuous level in the presence of cross-points. We show that its iterates can be uniquely decomposed into two parts, an even symmetric part that converges geometrically, like when there are no cross-points present, and an odd symmetric part, which generates a singularity at the cross-point and is not convergent. We illustrate our analysis with numerical experiments.

Keywords: domain decomposition methods, Dirichlet-Neumann methods, cross-points, elliptic problem

MSC Classification: 35J05 , 35D35 , 65N55 , 65N06

1 Introduction

Domain decomposition methods for partial differential equations are naturally defined and analyzed at the continuous level, like the original overlapping

Schwarz method from 1869 [1], and also the original Dirichlet-Neumann [2] and the Neumann-Neumann [3] method. Even FETI (Finite Element Tearing and Interconnect) was first presented at the continuous level in the original publication [4], as a minimization problem, before the authors proceeded to the finite element discretization that led to its name. For symmetric and positive definite problems, early domain decomposition research focused then however on condition number estimates for such methods used as preconditioners at the discrete level, leading to groundbreaking results, see e.g. [5] and references therein. The methods were thus intimately linked with the conjugate gradient method and not considered as standalone solvers, in contrast to multigrid methods for example [6, 7]. For more general problems which are non-symmetric and/or indefinite, condition numbers are not the key quantities anymore for understanding their convergence when used as preconditioners, and directly studying preconditioning properties for more general Krylov methods becomes difficult. There has therefore been an effort to also investigate the underlying iterative versions of these methods, and the study of their convergence properties at the continuous level, see for example [8] and references therein. This reveals many interesting properties of domain decomposition methods which are masked by the Krylov method that can correct convergence problems when the domain decomposition method is used as preconditioner. An interesting example is the Additive Schwarz method, which needs Krylov acceleration to be used, while Restricted Additive Schwarz does not, since it corresponds directly to the discretization of the parallel Schwarz method of Lions [9], and thus converges as a standalone iterative method [10, 11].

We are interested here in understanding the convergence properties of the Dirichlet-Neumann method in the presence of cross-points. Cross-points in domain decomposition methods have become a focus of attention over the past years because of an increasing interest in the domain decomposition research community to better understand the discretization of domain decomposition methods at cross-points, and the influence on the convergence of the iterative solution. For the Helmholtz equation and Després' seminal non-overlapping Schwarz method with Robin transmission conditions, cross-points do not hamper convergence [12], and the same holds more generally for non-overlapping optimized Schwarz methods when studied at the continuous level, see [13]. Care needs to be taken however when discretizing such methods, see for example [14–18], and this is even more important when higher order transmission conditions like Ventcell conditions are used [19–23]. For the Neumann-Neumann method, a well-posedness issue has been identified in the presence of cross-points, see [24], and the authors present a modification of the method to get around this difficulty. Also multitrace formulations pose problems at cross-points, and a solution involving specific non-local operators has been proposed in [25], see also [26] for a purely algebraic formulation. There was even a dedicated mini-symposium on cross-points in domain decomposition methods at the last international conference on domain decomposition methods, DD27, in Hong Kong [27].




Fig. 1: Domain Ω divided into four square subdomains colored in gray and white.

For the Dirichlet-Neumann method, the well-posedness issue in the presence of cross points was already mentioned in early work [28], but has so far not been fully analyzed for the iterative version of the algorithm. Most research works using this method or variants of this method do not encounter this problem. Indeed, many authors consider the case of domain decompositions with two subdomains [2, 29, 30] or many subdomains in stripes [31, 32], which excludes the presence of cross-points. Others use the Dirichlet-Neumann method as a preconditioner for a Krylov method, see for example [8, 33, 34], which hides the problematic behaviour of the Dirichlet-Neumann method. We focus here on a simple but instructive Laplace problem on a square divided into four squared subdomains of equal area. This allows us to develop a complete analytical understanding of the convergence of the Dirichlet-Neumann method in the presence of cross-points at the continuous level. We will show that the even symmetric part of the iteration converges like when no cross-points are present, but the odd symmetric part of the iteration generates singularities at cross-points and is thus not convergent. Even though our analysis is limited to the case of four subdomains, it reveals the behaviour of the iterates locally at the cross-point. Therefore it is representative of the behaviour of the iterates near cross-points for more general domain decompositions of grid-shape. The study on how to correct the convergence problem of the odd symmetric part will be addressed in a further research paper.

2 Geometry and model problem

As shown in Figure 1, we use as domain $\Omega \subset \mathbb{R}^2$ for our Laplace model problem the square $(-1, 1) \times (-1, 1)$, divided into four non-overlapping square subdomains Ω_i , $i \in \mathcal{I} := \{1, 2, 3, 4\}$, of equal area. With such a partition, there is one interior cross-point (red dot), and there are also four boundary cross-points (black dots). The presence of boundary cross-points has been identified as an obstacle for multitrace formulations, which requires some additional specific treatment, see for example [35, Section 2.4]. Conversely, these points are not an issue here for the Dirichlet-Neumann method. We denote the interfaces between adjacent subdomains by $\Gamma_{ij} := \text{int}(\partial\Omega_i \cap \partial\Omega_j)$, and the skeleton of

the partition by $\Gamma := \bigcup_{i,j} \bar{\Gamma}_{ij}$, and we have $\Gamma_{ij} = \Gamma_{ji}$ for all i, j . We further denote the interior of the intersection between $\partial\Omega_i$ and the boundary $\partial\Omega$ by $\partial\Omega_i^0 := \text{int}(\partial\Omega_i \cap \partial\Omega)$, and the interior of the left, right, bottom and top sides of Ω by $\partial\Omega_l, \partial\Omega_r, \partial\Omega_b, \partial\Omega_t$. Thus an arbitrary side of Ω is denoted by $\partial\Omega_\sigma$, where σ is in the set of indices $\mathcal{S} := \{l, r, b, t\}$. We use the same notation also for an arbitrary side of a subdomain Ω_i , namely $\partial\Omega_{i,\sigma}$ with $\sigma \in \mathcal{S}$.

2.1 Even and odd symmetric functions

We recall now the definition of an even/odd symmetric function in multivariate calculus, and a useful decomposition result, which we express here in terms of functions in L^p , for $p \in [1, +\infty]$.

Definition 1 (Symmetric set) Let $n \geq 1$ be an integer. A subset U of $\mathbb{R}^n$ is said to be *symmetric* if, for any $(x_1, \dots, x_n) \in \mathbb{R}^n$,

$$(x_1, \dots, x_n) \in U \implies (-x_1, \dots, -x_n) \in U.$$

Definition 2 (Even/odd symmetric functions) Let $U \subset \mathbb{R}^n$ be open and symmetric, and $p \in [1, +\infty]$. A function $h \in L^p(U)$ is called *even symmetric* if for almost all $(x_1, \dots, x_n) \in U$,

$$h(-x_1, \dots, -x_n) = h(x_1, \dots, x_n).$$

Similarly, the function h is called *odd symmetric* if for almost all $(x_1, \dots, x_n) \in U$,

$$h(-x_1, \dots, -x_n) = -h(x_1, \dots, x_n).$$

Theorem 1 (Even/odd decomposition) Let $U \subset \mathbb{R}^n$ be open and symmetric, and $p \in [1, +\infty]$. Every function $h \in L^p(U)$ can be uniquely decomposed as the sum of an even and an odd symmetric function, both in $L^p(U)$, which are called the *even symmetric part* and the *odd symmetric part* of the function. These functions, denoted by h_e and h_o , are for almost all $(x_1, \dots, x_n) \in U$ given by

$$\begin{aligned} h_e(x_1, \dots, x_n) &:= \frac{1}{2} (h(x_1, \dots, x_n) + h(-x_1, \dots, -x_n)), \\ h_o(x_1, \dots, x_n) &:= \frac{1}{2} (h(x_1, \dots, x_n) - h(-x_1, \dots, -x_n)). \end{aligned}$$

Proof Taking the sum, we see that $h_e + h_o = h$ almost everywhere (a.e.) in U . Then, regarding uniqueness, let us assume that there exists another couple of even/odd symmetric functions $(\tilde{h}_e, \tilde{h}_o) \neq (h_e, h_o)$ such that $\tilde{h}_e + \tilde{h}_o = h$ a.e. in U . It follows that

$$(h_e + h_o)(x_1, \dots, x_n) = (\tilde{h}_e + \tilde{h}_o)(x_1, \dots, x_n), \text{ for almost all } (x_1, \dots, x_n) \in U.$$

This implies that $h_e - \tilde{h}_e = \tilde{h}_o - h_o$ a.e. in U . Since the left hand-side is even symmetric and the right hand-side is odd symmetric, we must have $h_e - \tilde{h}_e = \tilde{h}_o - h_o = 0$ a.e. in U , which contradicts the initial assumption. $\square$

Definition 3 (Symmetry preservation of operators) Let X and Y be two Banach spaces such that $X \subset L^p(U)$ and $Y \subset L^q(V)$ for some open symmetric sets U, V and some $p, q \in [1, +\infty]$. An operator $T : X \rightarrow Y$ is said to *preserve symmetry* if it satisfies the following properties:

- for all $h \in X$ such that h is even symmetric, Th is even symmetric,
- for all $h \in X$ such that h is odd symmetric, Th is odd symmetric.

The Laplace operator $\Delta : H^2(\Omega) \rightarrow L^2(\Omega)$ preserves symmetry, which can be proved using the standard chain rule for C^2 functions together with the density of $C^\infty(\overline{\Omega})$ in $H^2(\Omega)$. In the same way, if p denotes an even symmetric function in $W^{\frac{1}{2},\infty}(\partial\Omega)$, one can prove that the operator $(\partial_n + p) : H^{\frac{3}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$ also preserves symmetry because for each $x \in \Gamma$, $n(-x) = -n(x)$.

2.2 Laplace model problem

We consider the Laplace problem with Dirichlet boundary condition on Ω , that is: find u solution to

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = g & \text{on } \partial\Omega, \end{cases} \quad (1)$$

where $f \in H^{-1}(\Omega)$ and $g \in H^{\frac{1}{2}}(\partial\Omega)$. Of course, since Ω is Lipschitz, it is known that (1) admits a unique solution $u \in H^1(\Omega)$. We also consider the Laplace problem with Robin boundary condition: find u solution to

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ (\partial_n + p)u = g & \text{on } \partial\Omega, \end{cases} \quad (2)$$

where $p \in L^\infty(\partial\Omega)$ ($p \geq 0$ a.e. on $\partial\Omega$) is even symmetric, $f \in H^{-1}(\Omega)$ and $g \in H^{-\frac{1}{2}}(\partial\Omega)$. To ensure that the problem is well-posed in $H^1(\Omega)$, we assume that p is strictly positive on a subset of $\partial\Omega$ of non-zero measure.

2.3 Regularity results

In this section, we briefly recall some important results about the theory of elliptic boundary value problems in nonsmooth two-dimensional domains, adapted to our specific context of a square. The general results in arbitrary polygons, together with their proofs, can be found in [36, Chapter 4], [37] and [38]. For a brief review on the subject, the reader is also referred to the lecture notes [39, Chapter 2].

$H^1(\Omega)$ *regularity*

As it has just been mentioned, the standard variational approach to study problems (1) and (2) leads to existence and uniqueness of solutions in $H^1(\Omega)$. However, for this result to hold, the regularity required on the boundary data is $g \in H^{\frac{1}{2}}(\partial\Omega)$ for the Dirichlet case and $g \in H^{-\frac{1}{2}}(\partial\Omega)$ for the Neumann (Robin)

case. This means that g can be identified as the trace on $\partial\Omega$ of a function in $H^1(\Omega)$ in the Dirichlet case. Similarly, in the Neumann (Robin) case, it can be identified as the normal derivative on $\partial\Omega$ of a function in $H^1(\Omega)$. When Ω is a polygon, the regularity on $\partial\Omega$ can be expressed by means of the regularity on each side of the polygon associated with so-called *compatibility relations* at the corners, see [37, Chapter 1]. Let us introduce the set of corners $\mathcal{C}$ which consists in four pairs of indices in $\mathcal{S}$:

$$\mathcal{C} := \{(l, b), (r, b), (r, t), (l, t)\} .$$

For the reader's convenience, the restriction of g to a part of the boundary $\partial\Omega_\sigma$ will be denoted by $g_\sigma := g|_{\partial\Omega_\sigma}$, for each $\sigma \in \mathcal{S}$. With these notations, we have the following useful characterization for the Dirichlet case: $g \in H^{\frac{1}{2}}(\partial\Omega)$ iff

$$g_\sigma \in H^{\frac{1}{2}}(\partial\Omega_\sigma), \forall \sigma \in \mathcal{S}, \text{ and } \int_0^\varepsilon |g_\sigma(s) - g_{\sigma'}(-s)|^2 \frac{ds}{s} < +\infty, \forall (\sigma, \sigma') \in \mathcal{C}, \quad (3)$$

for some small $\varepsilon > 0$. There exists a similar (and simpler) characterization for the Robin case: $g \in H^{-\frac{1}{2}}(\partial\Omega)$ iff

$$g_\sigma \in H^{-\frac{1}{2}}(\partial\Omega_\sigma), \forall \sigma \in \mathcal{S}. \quad (4)$$

Therefore, replacing the regularity conditions on the boundary data g in problems (1) and (2) by conditions (3) and (4) leads to well-posed formulations in $H^1(\Omega)$.

$H^2(\Omega)$ *regularity*

The existence and uniqueness results mentioned above only give us $H^1(\Omega)$ regularity for the solution, which means u might not even be continuous since $\Omega \subset \mathbb{R}^2$. Since we are interested in pointwise properties, especially what happens at the cross-point, we need more regular functions.

Theorem 2 (*Dirichlet case*) *If in addition to the previous assumptions, we have $f \in L^2(\Omega)$, $g_\sigma \in H^{\frac{3}{2}}(\partial\Omega_\sigma)$ for all $\sigma \in \mathcal{S}$, and $g_\sigma(0) = g_{\sigma'}(0)$ for all $(\sigma, \sigma') \in \mathcal{C}$, then the solution u to (1) is in $H^2(\Omega)$.*

Theorem 3 (*Robin case*) *If in addition to the previous assumptions, we have $f \in L^2(\Omega)$, $p|_{\partial\Omega_\sigma} \in C^\infty(\partial\Omega_\sigma)$ and $g_\sigma \in H^{\frac{1}{2}}(\partial\Omega_\sigma)$ for all $\sigma \in \mathcal{S}$, then the solution u to (2) is in $H^2(\Omega)$.*

We will also need regularity results for the mixed (Dirichlet/Neumann) problem for subproblems generated by the Dirichlet-Neumann method: let $\mathcal{D}$ and $\mathcal{N}$ be two subsets of $\mathcal{S}$ corresponding to the indices for Dirichlet and

Neumann boundary conditions. In order to avoid well-posedness issues, we assume that $\mathcal{D} \neq \emptyset$. The mixed problem reads: find u solution to

$$\begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = g_\sigma^\mathcal{D} & \text{on } \partial\Omega_\sigma, \text{ for } \sigma \in \mathcal{D}, \\ \partial_n u = g_\sigma^\mathcal{N} & \text{on } \partial\Omega_\sigma, \text{ for } \sigma \in \mathcal{N}. \end{cases} \quad (5)$$

The set of corners $\mathcal{C}$ can be split into three subsets $\mathcal{C} = \mathcal{C}_\mathcal{D} \cup \mathcal{C}_\mathcal{N} \cup \mathcal{C}_\mathcal{M}$ corresponding to Dirichlet corners, Neumann corners or mixed corners, such that for each $(\sigma, \sigma') \in \mathcal{C}$,

$$(\sigma, \sigma') \in \begin{cases} \mathcal{C}_\mathcal{D} & \text{if } \sigma, \sigma' \in \mathcal{D}, \\ \mathcal{C}_\mathcal{N} & \text{if } \sigma, \sigma' \in \mathcal{N}, \\ \mathcal{C}_\mathcal{M} & \text{if } \sigma \in \mathcal{D}, \sigma' \in \mathcal{N}. \end{cases}$$

Theorem 4 (*Dirichlet/Neumann case*) *If in addition to the previous assumptions, we have $f \in L^2(\Omega)$, $g_\sigma^\mathcal{D} \in H^{\frac{3}{2}}(\partial\Omega_\sigma)$ for $\sigma \in \mathcal{D}$, $g_\sigma^\mathcal{N} \in H^{\frac{1}{2}}(\partial\Omega_\sigma)$ for $\sigma \in \mathcal{N}$, and*

$$\begin{aligned} g_\sigma^\mathcal{D}(0) &= g_{\sigma'}^\mathcal{D}(0), \quad \forall (\sigma, \sigma') \in \mathcal{C}_\mathcal{D}, \\ \int_0^\varepsilon |g_\sigma^{\mathcal{D}'}(s) - g_{\sigma'}^\mathcal{N}(-s)|^2 \frac{ds}{s} &< +\infty, \quad \forall (\sigma, \sigma') \in \mathcal{C}_\mathcal{M}, \end{aligned}$$

then the solution u to (5) is in $H^2(\Omega)$.

Finally, let us state one last useful result about the regularity of such functions, which is a direct consequence of the Sobolev embedding theorem, see for example [40].

Proposition 5 *Let U be a bounded Lipschitz open subset of $\mathbb{R}^2$. Then, any function in $H^2(U)$ is also in $C^0(\overline{U})$.*

Proof From the Sobolev embedding theorem, since $2 > \frac{3}{2}$ and $U \subset \mathbb{R}^2$, we know that $H^2(U)$ is (continuously) embedded in the Hölder space $C_b^{0,\mu}(U)$ for some $\mu \in (0, 1)$. In addition, any function in $C_b^{0,\mu}(U)$ is uniformly continuous on U , therefore it can be extended to a continuous function on $\overline{U}$. $\square$

From now on, we will assume that the data satisfy the regularity required by the assumptions in Theorem 2 and Theorem 3. This ensures that the regularity of the solutions we deal with is at least $H^2(\Omega)$.

2.4 Even/odd symmetric decomposition

We now decompose the problems (1) and (2) into two subproblems, in order to analyze the subproblems separately. With Theorem 1, we can uniquely

decompose the data, and thus the associated problem, into the *even symmetric part* and the *odd symmetric part*. For problem (1), this leads to: find u^e and u^o solutions to

$$\begin{cases} -\Delta u^e = f_e & \text{in } \Omega, \\ u^e = g_e & \text{on } \partial\Omega, \end{cases} \quad (6)$$

$$\begin{cases} -\Delta u^o = f_o & \text{in } \Omega, \\ u^o = g_o & \text{on } \partial\Omega. \end{cases} \quad (7)$$

These subproblems are still well-posed and the solutions still have $H^2(\Omega)$ regularity. Using the symmetry preserving property of the operator $\Delta : H^2(\Omega) \rightarrow L^2(\Omega)$, we see that u^e is even symmetric and u^o is odd symmetric. Moreover, since $(-\Delta)$ is linear and the boundary conditions are linear as well, $u^e + u^o$ solves (1). Therefore $u^e + u^o = u$, and since the decomposition is unique, the solution to (6) u^e coincides with u_e , the even symmetric part of u , and similarly $u^o = u_o$. As we will see, the convergence analysis of the Dirichlet-Neumann method reveals different behaviors for the even and odd symmetric subproblems. Note that the geometric domain decomposition itself is also symmetric with respect to the origin $(0, 0)$.

As for the Dirichlet problem, we define the even and odd symmetric parts of (2), and still denote by u^e and u^o their solutions. When p is even symmetric, in addition to the operator Δ , we have seen that the operator $(\partial_n + p) : H^{\frac{3}{2}}(\partial\Omega) \rightarrow H^{\frac{1}{2}}(\partial\Omega)$ also preserves symmetry. Hence the even and odd symmetry properties of the solutions u^e and u^o . Thus, the linearity of (2) yields $u^e + u^o = u$, and the uniqueness of the decomposition finally leads to $u^e = u_e$ and $u^o = u_o$ like in the Dirichlet case.

3 Analysis of the Dirichlet-Neumann method

We use as in [8, Section 1.4] a gray and white coloring, see Figure 1, and define the sets of indices $\mathcal{I}_G := \{1 \leq i \leq 4 : \Omega_i \text{ is gray}\} = \{2, 4\}$ and $\mathcal{I}_W := \mathcal{I} \setminus \mathcal{I}_G = \{1, 3\}$. The transmission conditions are indicated in Figure 1, where "D" stands for Dirichlet and "N" for Neumann. Given an initial guess u^0 and a relaxation parameter $\theta \in \mathbb{R}$, each iteration $k \geq 1$ of the method can be split into two steps:

- **(Dirichlet step)** Solve for all $i \in \mathcal{I}_W$

$$\begin{cases} -\Delta u_i^k = f & \text{in } \Omega_i, \\ u_i^k = g & \text{on } \partial\Omega_i^0, \\ u_i^k = \theta u_j^{k-1} + (1 - \theta) u_i^{k-1} & \text{on } \Gamma_{ij}, \forall j \in \mathcal{I}_G \text{ s.t. } \Gamma_{ij} \neq \emptyset. \end{cases}$$

- **(Neumann step)** Solve for all $i \in \mathcal{I}_G$

$$\begin{cases} -\Delta u_i^k = f & \text{in } \Omega_i, \\ u_i^k = g & \text{on } \partial\Omega_i^0, \\ \partial_{n_i} u_i^k = -\partial_{n_j} u_j^k & \text{on } \Gamma_{ij}, \forall j \in \mathcal{I}_W \text{ s.t. } \Gamma_{ij} \neq \emptyset. \end{cases}$$

To start the Dirichlet-Neumann method, we need an initial guess u^0 , or equivalently $\lambda^0 := u^0|_{\Gamma}$, since only the traces of u^0 on the interfaces Γ_{ij} are used in the initialization step $k = 1$. This initial guess needs to satisfy the following compatibility condition.

Definition 4 (Compatible initial guess) An initial guess u^0 (or equivalently λ^0) is said to be compatible with the Dirichlet boundary condition if it satisfies $u^0|_{\partial\Omega \cap \Gamma} = g|_{\Gamma}$ (or equivalently $\lambda^0 = g$ a.e. on $\partial\Omega \cap \Gamma$ representing the set of boundary cross-points).

In what follows, we fix an initial guess λ^0 such that $\lambda^0 \in C^0(\Gamma) \cap H^{\frac{3}{2}}(\Gamma_{ij})$ for all (i, j) such that $\Gamma_{ij} \neq \emptyset$, and λ^0 is compatible with the boundary condition.

3.1 Case of the even symmetric part

We begin with applying the Dirichlet-Neumann method to the even symmetric part of problem (1).

Theorem 6 Taking λ_e^0 as the initial guess for the Dirichlet-Neumann method applied to the even symmetric part of problem (1) produces a sequence $\{u_e^k\}_k$ that converges geometrically to the solution u_e in the L^2 -norm and the broken H^1 -norm, for any $\theta \in (0, 1)$. Moreover, the convergence factor is given by $|1 - 2\theta|$, which also proves that this method becomes a direct solver for the specific choice $\theta = \frac{1}{2}$ ¹.

Proof We perform the first two iterations of the Dirichlet-Neumann method in terms of the local errors $e_i^k := u_i - u_i^k$ where $u_i := u|_{\Omega_i}$ is the restriction of the original solution to the i -th subdomain, see Section 4 (Example 1) for a numerical illustration.

Iteration $k = 1$, Dirichlet step: In Ω_1 , we solve

$$\begin{cases} -\Delta e_{e,1}^1 = 0 & \text{in } \Omega_1, \\ e_{e,1}^1 = 0 & \text{on } \partial\Omega_1^0, \\ e_{e,1}^1 = u_e - \lambda_e^0 & \text{on } \Gamma_{12} \cup \Gamma_{41}. \end{cases}$$

Since λ_e^0 is compatible with the even part of the Dirichlet boundary condition and $u_e|_{\partial\Omega_{1,\sigma}} \in H^{\frac{3}{2}}(\partial\Omega_{1,\sigma})$ for all $\sigma \in \mathcal{S}$, by Theorem 2, $e_{e,1}^1 \in H^2(\Omega_1)$ exists and is

¹The same result was already proved in [2], in the case of a symmetric decomposition with two subdomains without any restriction on the even/odd symmetric nature of the data.

unique. Moreover, we know that it is also continuous in $\overline{\Omega}_1$ due to Proposition 5. In the same way, we solve in Ω_3

$$\begin{cases} -\Delta e_{e,3}^1 = 0 & \text{in } \Omega_3, \\ e_{e,3}^1 = 0 & \text{on } \partial\Omega_3^0, \\ e_{e,3}^1 = u_e - \lambda_e^0 & \text{on } \Gamma_{23} \cup \Gamma_{34}. \end{cases}$$

Since $u_e - \lambda_e^0$ is even symmetric, it follows that the only solution $e_{e,3}^1 \in H^2(\Omega_3)$ to this problem verifies $e_{e,3}^1(x, y) = e_{e,1}^1(-x, -y)$, for all $(x, y) \in \overline{\Omega}_3$.

Iteration $k = 1$, Neumann step: Now, in Ω_2 , we solve

$$\begin{cases} -\Delta e_{e,2}^1 = 0 & \text{in } \Omega_2, \\ e_{e,2}^1 = 0 & \text{on } \partial\Omega_2^0, \\ \partial_x e_{e,2}^1 = \partial_x e_{e,1}^1 = -\partial_x(e_{e,1}^1(-x, y))|_{x=0} & \text{on } \Gamma_{12}, \\ \partial_y e_{e,2}^1 = \partial_y e_{e,3}^1 = \partial_y(e_{e,1}^1(-x, -y))|_{y=0} = -\partial_y(e_{e,1}^1(-x, y))|_{y=0} & \text{on } \Gamma_{23}. \end{cases}$$

This problem is well-posed in $H^1(\Omega_2)$. In addition, it is clear that the function defined in Ω_2 by $(x, y) \mapsto -e_{e,1}^1(-x, y)$ solves the problem. By uniqueness, one deduces that the solution $e_{e,2}^1$ is given by $e_{e,2}^1(x, y) = -e_{e,1}^1(-x, y)$, for all $(x, y) \in \Omega_2$. Note that this equality extends to the whole $\overline{\Omega}_2$. Again, in the exact same way, we get that the solution $e_{e,4}^1$ in Ω_4 is given by $e_{e,4}^1(x, y) = -e_{e,1}^1(x, -y)$, for all $(x, y) \in \overline{\Omega}_4$.

We are left with a recombined error e_e^1 (defined in $\Omega \setminus \Gamma$) that is discontinuous across all parts of the skeleton Γ where $e_{e,1}^1 \neq 0$, see Figure 3c. This may lead to discontinuities for the recombined solution $u_e^1 = u_e + e_e^1$. Also note that e_e^1 is even symmetric in $\Omega \setminus \Gamma$.

Iteration $k = 2$, Dirichlet step: In Ω_1 , we solve

$$\begin{cases} -\Delta e_{e,1}^2 = 0 & \text{in } \Omega_1, \\ e_{e,1}^2 = 0 & \text{on } \partial\Omega_1^0, \\ e_{e,1}^2 = \theta e_{e,2}^1 + (1 - \theta)e_{e,1}^1 = (1 - 2\theta)e_{e,1}^1 & \text{on } \Gamma_{12}, \\ e_{e,1}^2 = (1 - 2\theta)e_{e,1}^1 & \text{on } \Gamma_{41}. \end{cases}$$

The unique solution to this problem is $e_{e,1}^2 = (1 - 2\theta)e_{e,1}^1$. In the exact same way, we get in Ω_3 , $e_{e,3}^2 = (1 - 2\theta)e_{e,3}^1$.

Iteration $k = 2$, Neumann step: Now, in Ω_2 , we solve

$$\begin{cases} -\Delta e_{e,2}^2 = 0 & \text{in } \Omega_2, \\ e_{e,2}^2 = 0 & \text{on } \partial\Omega_2^0, \\ \partial_x e_{e,2}^2 = \partial_x e_{e,1}^2 = (1 - 2\theta)\partial_x e_{e,1}^1 & \text{on } \Gamma_{12}, \\ \partial_y e_{e,2}^2 = \partial_y e_{e,3}^2 = (1 - 2\theta)\partial_y e_{e,3}^1 & \text{on } \Gamma_{23}. \end{cases}$$

The unique solution to this problem is $e_{e,2}^2 = (1 - 2\theta)e_{e,2}^1$. Again, in the exact same way, we get in Ω_4 , $e_{e,4}^2 = (1 - 2\theta)e_{e,4}^1$. For all $\theta \in (0, 1)$, the recombined solution e_e^2 is exactly $(1 - 2\theta)e_e^1$.

Iterations $k \geq 3$: From the analysis of the first two iterations, it follows by induction that, at iteration k , one has

$$e_e^k = (1 - 2\theta)^{k-1} e_e^1 \quad \text{in } \Omega \setminus \Gamma,$$

see Figure 3d. Therefore, the proposed domain decomposition method converges geometrically to the solution u_e both in the L^2 -norm and the broken H^1 -norm for all $\theta \in (0, 1)$,

$$\|u_e - u_e^k\|_{L^2(\Omega)} \leq C |1 - 2\theta|^{k-1} \quad \text{and} \quad \sum_{i \in \mathcal{I}} \|u_{e,i} - u_{e,i}^k\|_{H^1(\Omega_i)} \leq C' |1 - 2\theta|^{k-1}.$$

Note that the recombined solution u_e^k is in general not continuous across the skeleton Γ . $\square$

Remark 1 It is actually not difficult to find an initial guess λ^0 satisfying the assumptions of Theorem 6: the simplest way is to build a piecewise linear function on Γ . If we denote P_k , $k \in \{1, \dots, 4\}$, the four boundary cross points, and set $\lambda^0(P_k) := g(P_k)$ for each k , then λ^0 is compatible with the Dirichlet boundary condition. Next, we set $\lambda^0(0, 0) := \frac{1}{4} \sum_k \lambda^0(P_k)$ and perform a linear reconstruction on each interface Γ_{ij} , which ensures that $\lambda^0 \in C^0(\Gamma)$.

Of course, any other choice for $\lambda^0(0, 0)$ would work as well, but this specific choice leads to a function λ^0 with minimal slopes as it satisfies

$$\frac{1}{4} \sum_k \lambda^0(P_k) = \arg \min_{\alpha} \sum_k |\lambda^0(P_k) - \alpha|^2.$$

3.2 Case of the odd symmetric part


Let us now turn to the odd symmetric part of problem (1). Given our initial guess, we obtain the following result.

Theorem 7 *The Dirichlet-Neumann method applied to the odd symmetric part of problem (1) is not well-posed. More specifically, taking λ_o^0 as the initial guess, there exists an integer $k_0 > 0$ such that the solution to the problem obtained at the k_0 -th iteration is not unique. In addition, all possible solutions $u_o^{k_0}$ are singular at the cross-point, with a leading singularity of type $(\ln r)^2$.*

The previous result shows that in this case, at some point, the Dirichlet-Neumann method is not applicable anymore. In practice, since we are able to find a solution to this ill-posed problem, one may wonder if it is possible to recover a nice behaviour of the method if we go past this iteration k_0 . The next theorem provides a negative answer.

Theorem 8 *If we let the Dirichlet-Neumann method go beyond the ill-posed iteration k_0 from Theorem 7, we end up with a sequence $\{u_o^k\}_{k \geq k_0}$ of non-unique iterates. Moreover, for each $k \geq k_0$, all possible solutions u_o^k are singular at the cross-point, with a leading singularity of type $(\ln r)^{2(k-k_0)+2}$.*

Before giving the proofs of these theorems, we need four technical lemmas which provide solutions to the Laplace problem on the two-dimensional cone

**Fig. 2:** Cone C and its boundaries.

$C := \mathbb{R}_+^* \times (-\frac{\pi}{4}, \frac{\pi}{4})$ with different types of Dirichlet and Neumann boundary conditions on $\partial C_- := \mathbb{R}_+^* \times \{-\frac{\pi}{4}\}$ and $\partial C_+ := \mathbb{R}_+^* \times \{\frac{\pi}{4}\}$, see Figure 2.

Lemma 9 *For any integer $q \geq 0$, there exist coefficients $(\alpha_{D,j,m}^q)_{j,m}$ such that the function v_D^q defined in polar coordinates for all $(r, \phi) \in C$ by*

$$v_D^q(r, \phi) := (\ln r)^q + \sum_{j,m=1}^{\lfloor \frac{q}{2} \rfloor} \alpha_{D,j,m}^q \phi^{2j} (\ln r)^{q-2m}$$

is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$ and solves

$$\begin{cases} -\Delta v = 0 & \text{in } C, \\ v = (\ln r)^q & \text{on } \partial C_- \cup \partial C_+. \end{cases} \quad (8)$$

Proof Let us proceed by induction.

Base case. The cases $q = 0$ and $q = 1$ are easily verified. Indeed, $v_D^0 := 1$ and $v_D^1 := \ln r$ clearly solve (8) for $q = 0$ and $q = 1$, respectively.

Induction step. Let $q \geq 1$ be fixed. We assume that the result of the lemma holds for all $\ell \leq q$. In order to find a solution to (8) for $q + 1$, we begin with the initial guess $(\ln r)^{q+1}$, which obviously satisfies the boundary condition. First, we compute a correction r_D^{q+1} such that $(\ln r)^{q+1} + r_D^{q+1}$ is harmonic in C . Then, since the boundary conditions are no longer satisfied due to this first correction, we compute a second correction $\tilde{r}_D^{q+1}$ such that $(\ln r)^{q+1} + r_D^{q+1} + \tilde{r}_D^{q+1}$ solves (8).

For the first step, we start by computing the Laplacian in polar coordinates. Let us recall its expression for some smooth function f depending on (r, ϕ)

$$\Delta f = \frac{\partial^2 f}{\partial r^2} + \frac{1}{r} \frac{\partial f}{\partial r} + \frac{1}{r^2} \frac{\partial^2 f}{\partial \phi^2}.$$

For our initial guess $(\ln r)^{q+1}$, this leads to

$$\Delta \left((\ln r)^{q+1} \right) = (q+1)q \frac{(\ln r)^{q-1}}{r^2}.$$

We can now cancel the term remaining on the right by adding on the left

$$\Delta \left((\ln r)^{q+1} - \frac{1}{2} \phi^2(q+1)q(\ln r)^{q-1} \right) = -\frac{1}{2} \phi^2(q+1) \cdots (q-2) \frac{(\ln r)^{q-3}}{r^2}.$$

Again, to remove the new term on the right, we add on the left

$$\begin{aligned} \Delta \left((\ln r)^{q+1} - \frac{1}{2} \phi^2(q+1)q(\ln r)^{q-1} + \frac{1}{4!} \phi^4(q+1) \cdots (q-2)(\ln r)^{q-3} \right) \\ = \frac{1}{4!} \phi^4(q+1) \cdots (q-4) \frac{(\ln r)^{q-5}}{r^2}, \end{aligned}$$

and continuing until the exponent on the $(\ln r)$ term in the right hand side reaches 0 or 1 gives

$$\Delta \left((\ln r)^{q+1} + \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} (-1)^j \frac{1}{(2j)!} \phi^{2j}(q+1) \cdots (q+2-2j)(\ln r)^{q+1-2j} \right) = 0.$$

Therefore, introducing $\beta_j^{q+1} := (-1)^j \binom{q+1}{2j}$ for all j , and defining the correction

$$r_D^{q+1} := \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \phi^{2j}(\ln r)^{q+1-2j},$$

we get a function $(\ln r)^{q+1} + r_D^{q+1}$ that is harmonic and is in $L^2(C) \cap C^\infty(\bar{C} \setminus \{(0,0)\})$.

For the second step, let us note that this function verifies

$$(\ln r)^{q+1} + r_D^{q+1} = (\ln r)^{q+1} + \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j} (\ln r)^{q+1-2j}$$

on $\partial C_- \cup \partial C_+$. From the induction hypothesis, we know that for each j , there exists a function v_D^{q+1-2j} that solves (8) for $q+1-2j$. Therefore, if we define

$$\tilde{r}_D^{q+1} := - \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j} v_D^{q+1-2j},$$

we obtain a function $v_D^{q+1} := (\ln r)^{q+1} + r_D^{q+1} + \tilde{r}_D^{q+1}$ that solves (8) for $q+1$ and is in $L^2(C) \cap C^\infty(\bar{C} \setminus \{(0,0)\})$. In addition, given the expressions of the corrections r_D^{q+1} and $\tilde{r}_D^{q+1}$, it follows that v_D^{q+1} can be written as

$$v_D^{q+1}(r, \phi) = (\ln r)^{q+1} + \sum_{j,m=1}^{\lfloor \frac{q+1}{2} \rfloor} \alpha_{D,j,m}^{q+1} \phi^{2j} (\ln r)^{q+1-2m},$$

where the coefficients $\alpha_{D,j,m}^{q+1}$ depend on the β_j^{q+1} and the $\alpha_{D,j,m}^\ell$, for $\ell \leq q-1$. $\square$

Lemma 10 *For any integer $q \geq 0$, there exist coefficients $(\alpha_{D',j,m}^q)_{j,m}$ such that the function $v_{D'}^q$, defined in polar coordinates for all $(r, \phi) \in C$ by*

$$v_{D'}^q(r, \phi) := \frac{4}{\pi} \phi (\ln r)^q + \sum_{j,m=1}^{\lfloor \frac{q}{2} \rfloor} \alpha_{D',j,m}^q \phi^{2j+1} (\ln r)^{q-2m}$$

is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$ and solves

$$\begin{cases} -\Delta v = 0 & \text{in } C, \\ v = -(\ln r)^q & \text{on } \partial C_-, \\ v = (\ln r)^q & \text{on } \partial C_+. \end{cases} \quad (9)$$

Proof Let us proceed by induction, in the same way as in the proof of Lemma 9.

Base case. It is not difficult to check that $v_{D'}^0 := \frac{4}{\pi}\phi$ and $v_{D'}^1 := \frac{4}{\pi}\phi \ln r$ solve (9) for $q = 0$ and $q = 1$, respectively.

Induction step. Let $q \geq 1$ be fixed. We assume that the result of the lemma holds for all $\ell \leq q$. As for Lemma 9, in order to find a solution to (9) for $q + 1$, we proceed in two steps, this time starting from $\frac{4}{\pi}\phi(\ln r)^{q+1}$ as initial guess. For the first step, following a similar iterative approach, we obtain a correction

$$r_{D'}^{q+1} := \frac{4}{\pi}\phi r_D^{q+1} = \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \phi^{2j+1} (\ln r)^{q+1-2j},$$

where $\beta_j^{q+1} := \frac{4}{\pi}(-1)^j \binom{q+1}{2j}$ for all j . This enables us to get a function $\frac{4}{\pi}\phi(\ln r)^{q+1} + r_{D'}^{q+1}$ that is harmonic and is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$. However, we have

$$\frac{4}{\pi}\phi(\ln r)^{q+1} + r_{D'}^{q+1} = \pm \left((\ln r)^{q+1} + \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \left(\frac{\pi}{4}\right)^{2j+1} (\ln r)^{q+1-2j} \right)$$

on $\partial C_\pm$. This notation means that the sign depends on which part of the boundary is considered: $+$ on ∂C_+ and $-$ on ∂C_- . Therefore, using the induction hypothesis, we introduce a second correction

$$\tilde{r}_{D'}^{q+1} := - \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \left(\frac{\pi}{4}\right)^{2j+1} v_{D'}^{q+1-2j},$$

such that $v_{D'}^{q+1} := \frac{4}{\pi}\phi(\ln r)^{q+1} + r_{D'}^{q+1} + \tilde{r}_{D'}^{q+1}$ solves (9) for $q + 1$ and is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$. In addition, this function can be written as

$$v_{D'}^{q+1}(r, \phi) = \frac{4}{\pi}\phi(\ln r)^{q+1} + \sum_{j,m=1}^{\lfloor \frac{q+1}{2} \rfloor} \alpha_{D',j,m}^{q+1} \phi^{2j+1} (\ln r)^{q+1-2m},$$

where the coefficients $\alpha_{D',j,m}^{q+1}$ depend on the β_j^{q+1} and the $\alpha_{D',j,m}^\ell$, for $\ell \leq q - 1$. $\square$

Lemma 11 *For any integer $q \geq 0$, there exist coefficients $(\alpha_{N,j,m}^q)_{j,m}$ such that the function v_N^q defined in polar coordinates for all $(r, \phi) \in C$ by*

$$v_N^q(r, \phi) := \phi(\ln r)^q + \sum_{j,m=1}^{\lfloor \frac{q}{2} \rfloor} \alpha_{N,j,m}^q \phi^{2j+1} (\ln r)^{q-2m}$$

is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$ and solves

$$\begin{cases} -\Delta v = 0 & \text{in } C, \\ \partial_\phi v = (\ln r)^q & \text{on } \partial C_- \cup \partial C_+. \end{cases} \quad (10)$$

Proof Again, we proceed by induction, in the same spirit as in the proofs of the previous lemmas.

Base case. Obviously, $v_N^0 := \phi$ and $v_N^1 := \phi \ln r$ solve (10) for $q = 0$ and $q = 1$, respectively.

Induction step. Let $q \geq 1$ be fixed. The result of the lemma is assumed to hold for all $\ell \leq q$. As before, we aim at finding a solution to (10) for $q + 1$ by proceeding in two steps, this time starting from $\phi(\ln r)^{q+1}$ as initial guess.

This initial guess is the same as in the proof of Lemma 10 (up to a factor $\frac{4}{\pi}$), therefore we have for the first correction

$$r_N^{q+1} := \phi r_D^{q+1} = \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} \beta_j^{q+1} \phi^{2j+1} (\ln r)^{q+1-2j},$$

where $\beta_j^{q+1} := (-1)^j \binom{q+1}{2j}$ for all j . This choice ensures that $\phi(\ln r)^{q+1} + r_N^{q+1}$ is harmonic and is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$. However, the boundary condition satisfied by this new function is

$$\partial_\phi \left(\phi(\ln r)^{q+1} + r_N^{q+1} \right) = (\ln r)^{q+1} + \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} (2j+1) \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j} (\ln r)^{q+1-2j}$$

on $\partial C_- \cup \partial C_+$. This time, the induction hypothesis leads to a second correction

$$\tilde{r}_N^{q+1} := - \sum_{j=1}^{\lfloor \frac{q+1}{2} \rfloor} (2j+1) \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j} v_N^{q+1-2j}.$$

Finally, we define $v_N^{q+1} := \phi(\ln r)^{q+1} + r_N^{q+1} + \tilde{r}_N^{q+1}$, which is solution to (10) for $q + 1$ and is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$. Moreover, we have for this function the expression

$$v_N^{q+1}(r, \phi) = \phi(\ln r)^{q+1} + \sum_{j,m=1}^{\lfloor \frac{q+1}{2} \rfloor} \alpha_{N,j,m}^{q+1} \phi^{2j+1} (\ln r)^{q+1-2m},$$

where the coefficients $\alpha_{N,j,m}^{q+1}$ depend on the β_j^{q+1} and the $\alpha_{N,j,m}^\ell$, for $\ell \leq q-1$. $\square$

Lemma 12 *For any integer $q \geq 0$, there exist coefficients $(\alpha_{N',j,m}^q)_{j,m}$ such that the function $v_{N'}^q$, defined in polar coordinates for all $(r, \phi) \in C$ by*

$$v_{N'}^q(r, \phi) := \frac{4}{\pi} \left(\frac{1}{2} \phi^2 (\ln r)^q - \frac{1}{(q+1)(q+2)} (\ln r)^{q+2} \right) + \sum_{j,m=2}^{\lfloor \frac{q}{2} \rfloor + 1} \alpha_{N',j,m}^q \phi^{2j} (\ln r)^{q+2-2m}$$

is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$ and solves

$$\begin{cases} -\Delta v = 0 & \text{in } C, \\ \partial_\phi v = -(\ln r)^q & \text{on } \partial C_-, \\ \partial_\phi v = (\ln r)^q & \text{on } \partial C_+. \end{cases} \quad (11)$$

Proof We keep proceeding by induction.

Base case. Take $q = 0$. Then we have $v_{N'}^0 := \frac{2}{\pi} \left(\phi^2 - (\ln r)^2 \right)$, which is indeed solution to (11) for $q = 0$.

Induction step. Let $q \geq 1$ be fixed. The result of the lemma is assumed to hold for all $\ell \leq q$. In order to find a solution to (11) for $q + 1$, we follow the usual two steps starting from the initial guess $\frac{2}{\pi} \phi^2 (\ln r)^{q+1}$, which satisfies the boundary conditions. For the first step, we reuse the computations performed in the proof of Lemma 9, replacing $q + 1$ by $q + 3$. Rewriting the function $(\ln r)^{q+3} + r_D^{q+3}$, we get that

$$(\ln r)^{q+3} - \frac{1}{2} \phi^2 (q+3)(q+2) (\ln r)^{q+1} + \sum_{j=2}^{\lfloor \frac{q+3}{2} \rfloor} (-1)^j \binom{q+3}{2j} \phi^{2j} (\ln r)^{q+3-2j}$$

is a harmonic function. We can easily deduce from this a first correction for our initial guess,

$$r_{N'}^{q+1} := -\frac{4}{\pi} \frac{1}{(q+3)(q+2)} (\ln r)^{q+3} + \sum_{j=2}^{\lfloor \frac{q+3}{2} \rfloor} \beta_j^{q+1} \phi^{2j} (\ln r)^{q+3-2j},$$

where $\beta_j^{q+1} := \frac{4}{\pi} \frac{(-1)^{j+1}}{(q+3)(q+2)} \binom{q+3}{2j}$ for all j . As desired, the new function $\frac{2}{\pi} \phi^2 (\ln r)^{q+1} + r_{N'}^{q+1}$ is harmonic and belongs to $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$. In addition, it satisfies

$$\partial_\phi \left(\frac{2}{\pi} \phi^2 (\ln r)^{q+1} + r_{N'}^{q+1} \right) = \pm \left((\ln r)^{q+1} + \sum_{j=2}^{\lfloor \frac{q+3}{2} \rfloor} (2j) \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j-1} (\ln r)^{q+3-2j} \right)$$

on $\partial C_\pm$. Again, using the induction hypothesis for all $\ell \leq q - 1$, we obtain a second correction

$$\tilde{r}_{N'}^{q+1} := - \sum_{j=2}^{\lfloor \frac{q+3}{2} \rfloor} (2j) \beta_j^{q+1} \left(\frac{\pi}{4} \right)^{2j-1} v_{N'}^{q+3-2j}.$$

Therefore, we are able to build a function $v_{N'}^{q+1} := \frac{2}{\pi} \phi^2 (\ln r)^{q+1} + r_{N'}^{q+1} + \tilde{r}_{N'}^{q+1}$ that solves (11) for $q + 1$ and is in $L^2(C) \cap C^\infty(\overline{C} \setminus \{(0,0)\})$, namely

$$\begin{aligned} v_{N'}^{q+1}(r, \phi) &= \frac{4}{\pi} \left(\frac{1}{2} \phi^2 (\ln r)^{q+1} - \frac{1}{(q+2)(q+3)} (\ln r)^{q+3} \right) \\ &\quad + \sum_{j,m=2}^{\lfloor \frac{q+1}{2} \rfloor + 1} \alpha_{N',j,m}^{q+1} \phi^{2j} (\ln r)^{q+3-2m}, \end{aligned}$$

where the coefficients $\alpha_{N',j,m}^{q+1}$ depend on the β_j^{q+1} and the $\alpha_{N',j,m}^\ell$, for $\ell \leq q - 1$. $\square$

We can now prove Theorem 7 and Theorem 8.

Proof (Theorem 7) We want to prove that the algorithm has a nice behavior up to some iteration k_0 , where the iterate becomes non-unique and singular near the cross-point, with a leading singularity of type $(\ln r)^2$. The idea is to show that, in general, after two iterations only, the approximate solution given by the method is singular near the cross-point. In order to do so, let us apply the Dirichlet-Neumann method step by step to (7), and write the local subproblems in terms of the local errors $e_{o,i}^k$.

Iteration $k = 1$, Dirichlet step: In Ω_1 , we solve

$$\begin{cases} -\Delta e_{o,1}^1 = 0 & \text{in } \Omega_1, \\ e_{o,1}^1 = 0 & \text{on } \partial\Omega_1^0, \\ e_{o,1}^1 = u_o - \lambda_o^0 & \text{on } \Gamma_{12} \cup \Gamma_{41}. \end{cases}$$

As for the even symmetric case, by Theorem 2, $e_{o,1}^1 \in H^2(\Omega_1) \subset C^0(\overline{\Omega}_1)$ exists and is unique. In the same way, we solve in Ω_3

$$\begin{cases} -\Delta e_{o,3}^1 = 0 & \text{in } \Omega_3, \\ e_{o,3}^1 = 0 & \text{on } \partial\Omega_3^0, \\ e_{o,3}^1 = u_o - \lambda_o^0 & \text{on } \Gamma_{23} \cup \Gamma_{34}. \end{cases}$$

Since $u_o - \lambda_o^0$ is odd symmetric, it follows that the unique solution $e_{o,3}^1$ to this problem verifies, for all $(x, y) \in \overline{\Omega}_3$, $e_{o,3}^1(x, y) = -e_{o,1}^1(-x, -y)$. Note that the recombined solution is continuous across the cross-point from Ω_1 to Ω_3 because $e_{o,1}^1(0, 0) = (u_o - \lambda_o^0)(0, 0) = 0$, since $(u_o - \lambda_o^0)$ is odd symmetric, see Figure 5b.

Iteration $k = 1$, Neumann step: Now, in Ω_2 , we solve

$$\begin{cases} -\Delta e_{o,2}^1 = 0 & \text{in } \Omega_2, \\ e_{o,2}^1 = 0 & \text{on } \partial\Omega_2^0, \\ \partial_x e_{o,2}^1 = \partial_x e_{o,1}^1 = -\partial_x(e_{o,1}^1(-x, y))|_{x=0} & \text{on } \Gamma_{12}, \\ \partial_y e_{o,2}^1 = \partial_y e_{o,3}^1 = -\partial_y(e_{o,1}^1(-x, -y))|_{y=0} = \partial_y(e_{o,1}^1(-x, y))|_{y=0} & \text{on } \Gamma_{23}. \end{cases}$$

Here again, we know the problem is well-posed in $H^1(\Omega_2)$. However, in contrast to the even symmetric case, we cannot argue that $\pm e_{o,1}^1(-x, y)$ solves this problem. There is a sign incompatibility in the boundary condition: minus sign on Γ_{12} and plus sign on Γ_{23} . The solution $e_{o,2}^1$ cannot be expressed explicitly. Nevertheless, since $e_{o,1}^1 \in H^2(\Omega_1)$, we know that the trace $e_{o,1}^1(\cdot, -1)$ on the bottom side of Ω_1 and the normal derivative $\partial_x e_{o,1}^1(0, \cdot)$ on the right side of Ω_1 satisfy the compatibility relation in Theorem 4 at the mixed corner $(0, -1)$. Therefore, as $e_{o,1}^1(\cdot, -1) = 0$, this compatibility relation is also satisfied by the boundary conditions enforced in the previous problem at this same corner $(0, -1)$ in Ω_2 . The same argument can be used for the other mixed corner $(1, 0)$. Moreover, the H^2 regularity of $e_{o,1}^1$ also implies that $\partial_x e_{o,1}^1 \in H^{\frac{1}{2}}(\Gamma_{12})$ and $\partial_y e_{o,3}^1 \in H^{\frac{1}{2}}(\Gamma_{23})$, which finally yields $e_{o,2}^1 \in H^2(\Omega_2) \subset C^0(\overline{\Omega}_2)$ by Theorem 4 and Proposition 5. Despite this additional regularity property, we are still not able to express the solution. Especially, we do not know the value of $e_{o,2}^1$ at $(0, 0)$. For the reader's convenience, let us denote it by $\delta^1 := e_{o,2}^1(0, 0)$. Then, in Ω_4 , we get that the solution $e_{o,4}^1$ is given by $e_{o,4}^1(x, y) = -e_{o,2}^1(-x, -y)$, for all $(x, y) \in \overline{\Omega}_4$. Therefore, the recombined solution jumps across the cross-point from δ^1 in Ω_2 to $-\delta^1$ in Ω_4 . This time, we are left with a recombined error e_o^1 that is odd symmetric in $\Omega \setminus \Gamma$, and discontinuous across the cross point, see Figure 5b.

Iteration $k = 2$, Dirichlet step: In Ω_1 , we solve

$$\begin{cases} -\Delta e_{o,1}^2 = 0 & \text{in } \Omega_1, \\ e_{o,1}^2 = 0 & \text{on } \partial\Omega_1^0, \\ e_{o,1}^2 = \theta e_{o,2}^1 + (1 - \theta)e_{o,1}^1 & \text{on } \Gamma_{12}, \\ e_{o,1}^2 = \theta e_{o,4}^1 + (1 - \theta)e_{o,1}^1 & \text{on } \Gamma_{41}. \end{cases} \quad (12)$$

Since the boundary conditions are continuous on $\bar{\Gamma}_{12}$ and $\bar{\Gamma}_{41}$, we are able to compute their limits at the cross-point, which yields

$$\begin{aligned} \lim_{\substack{(x,y) \rightarrow (0,0) \\ (x,y) \in \Gamma_{12}}} \left(\theta e_{o,2}^1 + (1-\theta)e_{o,1}^1 \right) (x,y) &= \theta \delta^1, \\ \lim_{\substack{(x,y) \rightarrow (0,0) \\ (x,y) \in \Gamma_{41}}} \left(\theta e_{o,4}^1 + (1-\theta)e_{o,1}^1 \right) (x,y) &= -\theta \delta^1. \end{aligned}$$

In other words, whenever $\delta^1 \neq 0$, the method enforces a discontinuous Dirichlet boundary condition at this step, which may lead to a singular solution since the compatibility relation (3) is no longer satisfied. Especially, the problem is not necessarily well-posed so the Dirichlet-Neumann method might not even be valid in this case. Note that there is no other discontinuity enforced since the boundary conditions on Γ_{12} and Γ_{41} extend to 0 at $\bar{\Gamma}_{12} \cap \bar{\Gamma}_1$ and $\bar{\Gamma}_{41} \cap \bar{\Gamma}_1$.

Case $\delta^1 \neq 0$. In what follows, we prove existence and uniqueness of $e_{o,1}^2$, exhibiting the type of singularity induced by this nonsmooth boundary condition. In order to do so, we try to decompose $e_{o,1}^2$ as the sum of a regular part v_1^2 and a singular part w_1^2 , in the same spirit as in [37] for more regular problems. First, for each $i \in \mathcal{I}$, let us introduce the angle $\phi_i \in (0, 2\pi)$ such that the rotation R_i of angle $-\phi_i$, given in polar coordinates by

$$R_i : (r, \phi) \mapsto (r, \phi - \phi_i),$$

maps the quadrant containing Ω_i onto the cone C . More specifically, we have

$$\phi_1 := \frac{5\pi}{4}, \quad \phi_2 := \frac{7\pi}{4}, \quad \phi_3 := \frac{\pi}{4}, \quad \phi_4 := \frac{3\pi}{4}.$$

Using these notations, we define $w_1^2 := (\theta \delta^1) \cdot v_{D'}^0 \circ R_1$, whose expression in polar coordinates (r, ϕ) reads

$$w_1^2(r, \phi) = \theta \delta^1 \frac{4}{\pi} (\phi - \phi_1). \quad (13)$$

We know from Lemma 10 that w_1^2 is of class C^∞ in $(\mathbb{R}_+)^2 \setminus \{(0,0)\}$, and that it satisfies

$$\begin{cases} -\Delta w_1^2 = 0 & \text{in } \mathbb{R}_-^* \times \mathbb{R}_-^*, \\ w_1^2 = \theta \delta^1 & \text{on } \{0\} \times \mathbb{R}_-^*, \\ w_1^2 = -\theta \delta^1 & \text{on } \mathbb{R}_-^* \times \{0\}. \end{cases}$$

Then, since we would like $v_1^2 + w_1^2$ to solve (12), we must define v_1^2 such that

$$\begin{cases} -\Delta v_1^2 = 0 & \text{in } \Omega_1, \\ v_1^2 = -w_1^2 & \text{on } \partial\Omega_1^0, \\ v_1^2 = \theta e_{o,2}^1 + (1-\theta)e_{o,1}^1 - w_1^2 & \text{on } \Gamma_{12}, \\ v_1^2 = \theta e_{o,4}^1 + (1-\theta)e_{o,1}^1 - w_1^2 & \text{on } \Gamma_{41}. \end{cases}$$

Note that the Dirichlet boundary condition enforced here is in $C^0(\partial\Omega_1) \cap H^{\frac{1}{2}}(\partial\Omega_{1,\sigma})$ for all $\sigma \in \mathcal{S}$. Therefore $v_1^2 \in H^1(\Omega_1)$ exists and is unique. Therefore, we have built (in a unique way) a solution $v_1^2 + w_1^2$ to (12) which is in $L^2(\Omega_1) \setminus H^1(\Omega_1)$. Indeed, it is easy to show that w_1^2 is in $L^2(\Omega_1)$, but not in $H^1(\Omega_1)$ since

$$|\nabla w_1^2|^2 = \left(\theta \delta^1 \frac{4}{\pi} \right)^2 \frac{1}{r^2}.$$

Now, in order to conclude that $e_{o,1}^2 = v_1^2 + w_1^2$, we must have uniqueness of the solution to (12) in $L^2(\Omega_1)$. This uniqueness property is indeed guaranteed since we know from [37, Theorem 4.4.3.3] that the subspace of all solutions $z \in L^2(\Omega_1)$ to

$$\begin{cases} -\Delta z = 0 & \text{in } \Omega_1, \\ z = 0 & \text{on } \partial\Omega_1, \end{cases}$$

is of dimension 0. Hence $e_{o,1}^2 := v_1^2 + w_1^2$ exists and is unique.

In the same way, one can conclude that, in Ω_3 , $e_{o,3}^2 := v_3^2 + w_3^2$ exists and is unique, with $v_3^2 \in H^1(\Omega_3)$ and $w_3^2 \in L^2(\Omega_3) \setminus H^1(\Omega_3)$. Of course, v_3^2 and w_3^2 can be obtained immediately from v_1^2 and w_1^2 using symmetry arguments, which gives for w_3^2

$$w_3^2(r, \phi) = -\theta\delta^1 \frac{4}{\pi} (\phi - \phi_3). \quad (14)$$

It is now clear that the algorithm generates a singular solution at this step. In order to estimate how the singularity propagates in the Neumann step, let us keep going and see what happens.

Iteration $k = 2$, Neumann step: In Ω_2 , we solve

$$\begin{cases} -\Delta e_{o,2}^2 = 0 & \text{in } \Omega_2, \\ e_{o,2}^2 = 0 & \text{on } \partial\Omega_2^0, \\ \partial_x e_{o,2}^2 = \partial_x e_{o,1}^2 & \text{on } \Gamma_{12}, \\ \partial_y e_{o,2}^2 = \partial_y e_{o,3}^2 & \text{on } \Gamma_{23}. \end{cases} \quad (15)$$

Due to the lack of regularity of $e_{o,1}^2$ and $e_{o,3}^2$, standard results fail to apply, so that existence and uniqueness of $e_{o,2}^2$ are not guaranteed. As in subdomains Ω_1 and Ω_3 , we decompose $e_{o,2}^2$ as the sum of a regular part v_2^2 and a singular part w_2^2 , exhibiting the singularity of w_2^2 . Using the previous decompositions for $e_{o,1}^2$ and $e_{o,3}^2$, we rewrite the boundary conditions on Γ_{12} and Γ_{23} ,

$$\begin{cases} \partial_x e_{o,2}^2 = \partial_x v_1^2 + \partial_x w_1^2 & \text{on } \Gamma_{12}, \\ \partial_y e_{o,2}^2 = \partial_y v_3^2 + \partial_y w_3^2 & \text{on } \Gamma_{23}. \end{cases}$$

Then, let us introduce the function $w_2^2 := -(\theta\delta^1 \frac{4}{\pi}) \cdot v_N^0 \circ R_2$, given in polar coordinates by

$$w_2^2(r, \phi) = -\theta\delta^1 \frac{8}{\pi^2} \left[(\phi - \phi_2)^2 - (\ln r)^2 \right]. \quad (16)$$

From Lemma 12, we know that $w_2^2 \in C^\infty(\mathbb{R}_+ \times \mathbb{R}_- \setminus \{(0, 0)\})$, and that it satisfies

$$\begin{cases} -\Delta w_2^2 = 0 & \text{in } \mathbb{R}_+^* \times \mathbb{R}_-^*, \\ \partial_x w_2^2 = \partial_x w_1^2 = -\left(\theta\delta^1 \frac{4}{\pi}\right) \frac{1}{y} & \text{on } \{0\} \times \mathbb{R}_-^*, \\ \partial_y w_2^2 = \partial_y w_3^2 = -\left(\theta\delta^1 \frac{4}{\pi}\right) \frac{1}{x} & \text{on } \mathbb{R}_+^* \times \{0\}. \end{cases}$$

This time, in order for $v_2^2 + w_2^2$ to solve the problem for $e_{o,2}^2$, we define v_2^2 such that

$$\begin{cases} -\Delta v_2^2 = 0 & \text{in } \Omega_2, \\ v_2^2 = -w_2^2 & \text{on } \partial\Omega_2^0, \\ \partial_x v_2^2 = \partial_x v_1^2 & \text{on } \Gamma_{12}, \\ \partial_y v_2^2 = \partial_y v_3^2 & \text{on } \Gamma_{23}. \end{cases}$$

Given the regularities of w_2^2 , v_1^2 and v_3^2 , we deduce that v_2^2 exists and is unique in $H^1(\Omega_2)$. Again, we have built (in a unique way) a solution $v_2^2 + w_2^2$ to (15), where $v_2^2 \in H^1(\Omega_2)$ and $w_2^2 \in L^2(\Omega_2) \setminus H^1(\Omega_2)$. But this is not enough to conclude that $e_{o,2}^2 = v_2^2 + w_2^2$. As for Ω_1 , we know that this last equality holds provided that the subspace of all solutions $z \in L^2(\Omega_2)$ to

$$\begin{cases} -\Delta z = 0 & \text{in } \Omega_2, \\ z = 0 & \text{on } \partial\Omega_2^0, \\ \partial_n z = 0 & \text{on } \Gamma_{12} \cup \Gamma_{23}, \end{cases}$$

is of dimension 0. Unfortunately, it follows from [37, Theorem 4.4.3.3] that its dimension is 1, and that it is spanned by a function in $L^2(\Omega_2)$, say z_2 , that admits a singularity of type $\ln r$ at the cross-point. Therefore, one has that $e_{o,2}^2$ is not unique, and it can be written as $e_{o,2}^2 = v_2^2 + w_2^2 + C_2^2 z_2$, for some constant $C_2^2 \in \mathbb{R}$. However, no matter the value of C_2^2 , one may always deduce that, in a neighborhood $\mathcal{V}_0$ of $(0,0)$,

$$e_{o,2}^2 \simeq \theta \delta^1 \frac{8}{\pi^2} (\ln r)^2 \quad \text{in } \Omega_2 \cap \mathcal{V}_0. \quad (17)$$

Note that there are actually three singular terms in $e_{o,2}^2$: $(\phi - \phi_2)^2$, $(\ln r)^2$ and $\ln r$. So the leading singularity is indeed $(\ln r)^2$.

Besides, one gets a similar result for $e_{o,4}^2$. That is $e_{o,4}^2$ is not unique and can be written as $e_{o,4}^2 = v_4^2 + w_4^2 + C_4^2 z_4$ for some $C_4^2 \in \mathbb{R}$, where $v_4^2 \in H^1(\Omega_4)$, $w_4^2 \in L^2(\Omega_4) \setminus H^1(\Omega_4)$ is given by

$$w_4^2(r, \phi) = \theta \delta^1 \frac{8}{\pi^2} \left[(\phi - \phi_4)^2 - (\ln r)^2 \right], \quad (18)$$

and $z_4 \in L^2(\Omega_4)$ admits a singularity of type $\ln r$ near the cross-point. Of course, one obtains a similar asymptotic result in the neighborhood $\mathcal{V}_0$ of $(0,0)$, i.e.

$$e_{o,4}^2 \simeq -\theta \delta^1 \frac{8}{\pi^2} (\ln r)^2 \quad \text{in } \Omega_4 \cap \mathcal{V}_0.$$

Note that in this case, the integer k_0 in the statement of the theorem equals 2.

Case $\delta^1 = 0$. In this case, well-posedness is guaranteed and no singularity is generated by the method at the current iteration $k = 2$. Indeed, the method behaves exactly as in the first iteration and all $e_{o,i}^2$ are well defined. Thus, we can introduce $\delta^2 \in \mathbb{R}$ such that $e_{o,2}^2(0,0) = -e_{o,4}^2(0,0) =: \delta^2$. Then we are again facing two possible situations: $\delta^2 \neq 0$, in which case well-posedness is lost and a $(\ln r)^2$ singularity is generated at the next iteration $k = 3$ (i.e. $k_0 = 3$), or $\delta^2 = 0$, in which case we still have the same behavior as in the first iteration. If we keep going with the same reasoning, we end up with two possible cases: either there exists some integer $k_0 > 1$ such that all iterates are uniquely defined and regular up to k_0 and both regularity and well-posedness are lost at $k = k_0$, or $\delta^k = 0$ for all $k \geq 1$, in which case all iterates are well defined and regular. This last case, which has never been encountered in practice, is not treated here as it seems difficult to study convergence properties in this specific situation. □

Proof (Theorem 8) Here, we study how the singularity exhibited in Theorem 7 propagates through the next iterations $k \geq k_0$. To begin, we assume that the algorithm is capable of finding one of the solutions to a problem for which uniqueness is not

guaranteed (typically problem (15)). To simplify notations, we introduce the integer $p \geq 0$ such that $k = k_0 + p$. Then, we claim that at iteration k , for each $i \in \mathcal{I}$, there exists a regular function $v_i^k \in H^1(\Omega_i)$ and real coefficients $(\gamma_{i,j,m}^k)_{j,m}$ such that the local error $e_{o,i}^k$ is given by

$$e_{o,i}^k = v_i^k + \sum_{m=0}^{2p+1} \sum_{j=0}^{2p} \gamma_{i,j,m}^k (\phi - \phi_i)^m (\ln r)^j, \quad \text{if } i \in \mathcal{I}_W, \quad (19)$$

$$e_{o,i}^k = v_i^k + \sum_{m=0}^{2p+2} \sum_{j=0}^{2p+2} \gamma_{i,j,m}^k (\phi - \phi_i)^m (\ln r)^j, \quad \text{if } i \in \mathcal{I}_G. \quad (20)$$

In addition, for $i \in \mathcal{I}_W$, we have $\gamma_{i,2p,m}^k = 0$ if $m > 1$, which means that the leading singularity in Ω_i is of type $(\phi - \phi_i)(\ln r)^{2p}$. And for $i \in \mathcal{I}_G$, we have $\gamma_{i,2p+2,m}^k = 0$ if $m \neq 0$, thus the leading singularity in Ω_i is of type $(\ln r)^{2p+2}$.

To prove this, we proceed by induction and use the results of the four lemmas stated earlier.

Base case. The case $k = k_0$, or equivalently $p = 0$, has already been seen in the proof of Theorem 7. Indeed, we have shown that the Dirichlet step in Ω_i ($i \in \mathcal{I}_W$) led to a local error with a singular part $w_i^{k_0}$ that matches the one in (19) for $k = k_0$, see expression (13) or (14). In addition, the Neumann step in Ω_i ($i \in \mathcal{I}_G$) led to a local error with a singular part $w_i^{k_0} + C_i^{k_0} z_i$, which can be replaced by $w_i^{k_0} + C_i^{k_0} (\ln r)$ up to some changes in the regular part $v_i^{k_0}$. It follows from expressions (16) and (18) that this matches the singular part in (20) for $k = k_0$.

Induction step. Let $k > k_0$, or equivalently $p > 0$, be fixed. Assuming our statement holds for any integer $\ell \leq k$, let us prove that it still holds for $k + 1$. Given $e_{o,i}^k$ for each $i \in \mathcal{I}$, the only way to get information about $e_{o,i}^{k+1}$ is to perform the Dirichlet and Neumann steps of the algorithm.

Dirichlet step: In Ω_1 we solve

$$\begin{cases} -\Delta e_{o,1}^{k+1} = 0 & \text{in } \Omega_1, \\ e_{o,1}^{k+1} = 0 & \text{on } \partial\Omega_1^0, \\ e_{o,1}^{k+1} = \theta e_{o,2}^k + (1 - \theta) e_{o,1}^k & \text{on } \Gamma_{12}, \\ e_{o,1}^{k+1} = \theta e_{o,4}^k + (1 - \theta) e_{o,1}^k & \text{on } \Gamma_{41}. \end{cases} \quad (21)$$

As previously, we know from [37, Theorem 4.4.3.3] that if there exists a solution to (21) that is in $L^2(\Omega_1)$, then it is unique. In order to prove existence, we use the induction hypothesis for $e_{o,2}^k$ and $e_{o,4}^k$, then we decompose $e_{o,1}^{k+1} = \theta \tilde{w}_1^{k+1} + (1 - \theta) e_{o,1}^k$, where $\tilde{w}_1^{k+1}$ is for $1 \leq j \leq 2p + 2$ the sum of the solutions to the boundary value problems

$$\begin{aligned} (a) \quad & \begin{cases} v = v_2^k & \text{on } \Gamma_{12}, \\ v = v_4^k & \text{on } \Gamma_{41}. \end{cases} \\ (b)_j \quad & \begin{cases} v = \mu_{2,j}^k (\ln r)^j & \text{on } \Gamma_{12}, \\ v = -\mu_{2,j}^k (\ln r)^j & \text{on } \Gamma_{41}. \end{cases} \\ (c)_j \quad & \begin{cases} v = \nu_{2,j}^k (\ln r)^j & \text{on } \Gamma_{12}, \\ v = \nu_{2,j}^k (\ln r)^j & \text{on } \Gamma_{41}, \end{cases} \end{aligned} \quad \text{and} \quad \begin{cases} -\Delta v = 0 & \text{in } \Omega_1, \\ v = 0 & \text{on } \partial\Omega_1^0, \end{cases} \quad (22)$$

where we have introduced for each j

$$\mu_{2,j}^k := \sum_{n=0}^{p+1} \gamma_{2,j,2n}^k \left(\frac{\pi}{4}\right)^{2n} \quad \text{and} \quad \nu_{2,j}^k := - \sum_{n=0}^p \gamma_{2,j,2n+1}^k \left(\frac{\pi}{4}\right)^{2n+1}.$$

Note that we have performed some simplifications using the equality $\gamma_{2,j,m}^k = -\gamma_{4,j,m}^k$ which holds for all k, j, m , due to the odd symmetry of the problem. Now, we consider each part of (22), and express the general form of its solution. First, since v_2^k and v_4^k are smooth and verify $v_2^k(x, y) = -v_4^k(-x, -y)$ for almost all $(x, y) \in \Omega_2$, we have already seen (in problem (12)) that there exists a solution

$$\tilde{w}_a^{k+1} \in \text{span}\{(\phi - \phi_1)\} + H^1(\Omega_1)$$

to part (a), where the coefficient in the linear combination depends on the jump $v_2^k(0, 0)$ at the cross-point. Then, for each j , we get from Lemma 10 that there exists a solution

$$\begin{aligned} \tilde{w}_{b_j}^{k+1} &\in \mu_{2,j}^k \frac{4}{\pi} (\phi - \phi_1) (\ln r)^j \\ &+ \text{span}\{(\phi - \phi_1)^m (\ln r)^q \mid 3 \leq m \leq j+1, 0 \leq q \leq j-2\} + H^1(\Omega_1) \end{aligned}$$

to part (b)_j. In addition, we also get from Lemma 9 that there exists a solution

$$\tilde{w}_{c_j}^{k+1} \in \nu_{2,j}^k (\ln r)^j + \text{span}\{(\phi - \phi_1)^m (\ln r)^q \mid 2 \leq m \leq j, 0 \leq q \leq j-2\} + H^1(\Omega_1)$$

to part (c)_j. Summing up all these contributions over the values of j , we deduce that there exists a function $\tilde{v}_1^{k+1} \in H^1(\Omega_1)$ and coefficients $(\tilde{\gamma}_{1,j,m}^{k+1})_{j,m}$ such that

$$\tilde{w}_1^{k+1} = \tilde{v}_1^{k+1} + \sum_{m=0}^{2p+3} \sum_{j=0}^{2p+2} \tilde{\gamma}_{1,j,m}^{k+1} (\phi - \phi_i)^m (\ln r)^j$$

solves (22) in $L^2(\Omega_1)$, with $\tilde{\gamma}_{1,2p+2,m}^{k+1} = 0$ for $m > 1$. Finally, defining $e_{o,1}^{k+1} := \theta \tilde{w}_1^{k+1} + (1 - \theta) e_{o,1}^k$, we obtain a solution to (21) that is in $L^2(\Omega_1)$, which also proves that it is unique in $L^2(\Omega_1)$. Moreover, using the previous decomposition for $\tilde{w}_1^{k+1}$ and the induction hypothesis for $e_{o,1}^k$, we get that there exists a function $v_1^{k+1} := \theta \tilde{v}_1^{k+1} + (1 - \theta) v_1^k \in H^1(\Omega_1)$ and coefficients $(\gamma_{1,j,m}^{k+1})_{j,m}$ such that

$$e_{o,1}^{k+1} = v_1^{k+1} + \sum_{m=0}^{2p+3} \sum_{j=0}^{2p+2} \gamma_{1,j,m}^{k+1} (\phi - \phi_1)^m (\ln r)^j,$$

with $\gamma_{1,2p+2,m}^{k+1} = \theta \tilde{\gamma}_{1,2p+2,m}^{k+1} = 0$ for $m > 1$. Obviously, the same result (existence, uniqueness, and decomposition) holds for $e_{o,3}^{k+1}$ in Ω_3 due to the odd symmetry property of the problem.

Neumann step: In Ω_2 we solve

$$\begin{cases} -\Delta e_{o,2}^{k+1} = 0 & \text{in } \Omega_2, \\ e_{o,2}^{k+1} = 0 & \text{on } \partial\Omega_2^0, \\ \partial_x e_{o,2}^{k+1} = \partial_x e_{o,1}^{k+1} & \text{on } \Gamma_{12}, \\ \partial_y e_{o,2}^{k+1} = \partial_y e_{o,3}^{k+1} & \text{on } \Gamma_{23}. \end{cases} \quad (23)$$

We have already seen (again from [37, Theorem 4.4.3.3]) that, if there exists a solution to (23) that is in $L^2(\Omega_2)$, then the set of all solutions is an affine subspace of dimension 1. As in the Dirichlet step, in order to prove existence, we decompose $e_{o,2}^{k+1}$ using the decompositions obtained for $e_{o,1}^{k+1}$ and $e_{o,3}^{k+1}$. More specifically, we consider for $1 \leq j \leq 2p+2$ the sum of the solutions to the boundary value problems

$$\begin{aligned} & (a) \quad \begin{cases} \partial_\phi v = \partial_\phi v_1^{k+1} & \text{on } \Gamma_{12}, \\ \partial_\phi v = \partial_\phi v_3^{k+1} & \text{on } \Gamma_{23}. \end{cases} \\ & \begin{cases} -\Delta v = 0 & \text{in } \Omega_2, \\ v = 0 & \text{on } \partial\Omega_2^0, \end{cases} \quad \text{and} \quad (b)_j \quad \begin{cases} \partial_\phi v = \mu_{1,j}^{k+1} (\ln r)^j & \text{on } \Gamma_{12}, \\ \partial_\phi v = -\mu_{1,j}^{k+1} (\ln r)^j & \text{on } \Gamma_{23}. \end{cases} \\ & (c)_j \quad \begin{cases} \partial_\phi v = \nu_{1,j}^{k+1} (\ln r)^j & \text{on } \Gamma_{12}, \\ \partial_\phi v = \nu_{1,j}^{k+1} (\ln r)^j & \text{on } \Gamma_{23}, \end{cases} \end{aligned} \quad (24)$$

where we have introduced for each j

$$\mu_{1,j}^{k+1} := \sum_{n=0}^{p+1} \gamma_{1,j,2n+1}^{k+1} (2n+1) \left(\frac{\pi}{4}\right)^{2n} \quad \text{and} \quad \nu_{1,j}^{k+1} := \sum_{n=0}^p \gamma_{1,j,2n+2}^{k+1} (2n+2) \left(\frac{\pi}{4}\right)^{2n+1}.$$

This time we have used the equality $\gamma_{1,j,m}^{k+1} = -\gamma_{3,j,m}^{k+1}$ to simplify the formulations. Let us now study separately each part of (24), and express the general form of its solution. To begin with, due to the regularities of v_1^{k+1} and v_3^{k+1} , we know from standard results that there exists a (unique) solution

$$w_a^{k+1} \in H^1(\Omega_2)$$

to part (a). Then, for each j , we get from Lemma 12 that there exists a solution

$$\begin{aligned} w_{b_j}^{k+1} & \in \mu_{1,j}^{k+1} \frac{4}{\pi} \left(\frac{1}{2} (\phi - \phi_2)^2 (\ln r)^j - \frac{1}{(j+1)(j+2)} (\ln r)^{j+2} \right) \\ & + \text{span} \{ (\phi - \phi_2)^m (\ln r)^q \mid 4 \leq m \leq j+2, 0 \leq q \leq j-2 \} + H^1(\Omega_2) \end{aligned}$$

to part (b)_j. Finally, we get from Lemma 11 that there exists a solution

$$\begin{aligned} w_{c_j}^{k+1} & \in \nu_{1,j}^{k+1} (\phi - \phi_2) (\ln r)^j \\ & + \text{span} \{ (\phi - \phi_2)^m (\ln r)^q \mid 3 \leq m \leq j+1, 0 \leq q \leq j-2 \} + H^1(\Omega_2) \end{aligned}$$

to part (c)_j. Combining these results, we end up with a function

$$v_2^{k+1} + \sum_{m=0}^{2p+4} \sum_{j=0}^{2p+4} \gamma_{2,j,m}^{k+1} (\phi - \phi_2)^m (\ln r)^j$$

that solves (24) in $L^2(\Omega_2)$, where $v_2^{k+1} \in H^1(\Omega_2)$ and the coefficients $(\gamma_{2,j,m}^{k+1})_{j,m}$ satisfy $\gamma_{2,2p+4,m}^{k+1} = 0$ if $m \neq 0$. This means that every solution $e_{o,2}^{k+1}$ to (24) is given by the general expression

$$e_{o,2}^{k+1} = v_2^{k+1} + \sum_{m=0}^{2p+4} \sum_{j=0}^{2p+4} \gamma_{2,j,m}^{k+1} (\phi - \phi_2)^m (\ln r)^j + C_2^{k+1} z_2,$$

with $C_2^{k+1} \in \mathbb{R}$. Since the singularity in z_2 is of type $\ln r$, it follows that every solution can be written as

$$e_{o,2}^{k+1} = v_2^{k+1} + \sum_{m=0}^{2p+4} \sum_{j=0}^{2p+4} \gamma_{2,j,m}^{k+1} (\phi - \phi_2)^m (\ln r)^j,$$

up to some modification of the regular part v_2^{k+1} and the coefficient $\gamma_{2,1,0}^{k+1}$. Again, the same result can be deduced for $e_{o,4}^{k+1}$ in Ω_4 .

This ends the proof of our claim about the expression of the local errors $e_{o,i}^k$ (see (19) and (20)). The type of singularity generated by the domain decomposition method at iteration k is directly given by those expressions, so the proof of the result is complete. $\square$

Remark 2 Following the same computations as in the previous proof while taking care of the coefficient in front of the leading singularity at each step enables us to end up with the following approximations of the errors in subdomains $\Omega_1, \dots, \Omega_4$ at iteration $k = k_0 + p$

$$\begin{aligned} e_{o,i}^k &\simeq \pm \theta \delta^1 \frac{4}{\pi} (\phi - \phi_1) \frac{1}{(2p)!} \left(\theta \left(\frac{4}{\pi} \ln r \right)^2 \right)^p, & \text{for } i \in \mathcal{I}_W, \\ e_{o,i}^k &\simeq \pm \delta^1 \frac{1}{(2p+2)!} \left(\theta \left(\frac{4}{\pi} \ln r \right)^2 \right)^{p+1}, & \text{for } i \in \mathcal{I}_G, \end{aligned} \quad (25)$$

where the correct sign is $+$ for $i \in \{1, 2\}$ and $-$ for $i \in \{3, 4\}$.

The results given by Theorem 6, Theorem 7 and Theorem 8 can be easily extended to the Laplace problem with Robin boundary conditions. In this case, no compatibility condition is required, we only impose that the initial guess λ^0 satisfies the regularity assumption that $\lambda^0 \in C^0(\Gamma) \cap H^{\frac{3}{2}}(\Gamma_{ij})$ for all (i, j) such that $\Gamma_{ij} \neq \emptyset$.

Theorem 13 Taking λ_e^0 as the initial guess for the Dirichlet-Neumann method applied to the even symmetric part of problem (2) produces a sequence $\{u_e^k\}_k$ that converges geometrically to the solution u_e with respect to the L^2 -norm and the broken H^1 -norm. Moreover, the convergence factor is given by $|1 - 2\theta|$, which also proves that this method becomes a direct solver for the specific choice $\theta = \frac{1}{2}$.

Theorem 14 The Dirichlet-Neumann method applied to the odd symmetric part of problem (2) is not well-posed. More specifically, taking λ_o^0 as the initial guess, there exists an integer $k_0 > 0$ such that the solution to the problem obtained at the k_0 -th iteration is not unique. In addition, all possible solutions $u_o^{k_0}$ are singular at the cross-point, with a leading singularity of type $(\ln r)^2$.

Theorem 15 If we let the Dirichlet-Neumann method go beyond the ill-posed iteration k_0 from Theorem 14, we end up with a sequence $\{u_o^k\}_{k \geq k_0}$ of non-unique iterates. Moreover, for each $k \geq k_0$, all possible u_o^k are singular at the cross-point, with a leading singularity of type $(\ln r)^{2(k-k_0)+2}$.

Proof The proofs can be obtained by following the same steps as in the proofs of Theorem 6, Theorem 7 and Theorem 8. $\square$

Remark 3 Note that the formulas given by the asymptotic analysis near the origin remain valid in the Robin case. Indeed, as this study has been conducted in the neighborhood of $(0, 0)$, what happens "far away" from this point (e.g. on the boundary $\partial\Omega$) has no influence on the results.

4 Numerical experiments

We now illustrate the theoretical results obtained in the previous section. The square domain Ω is discretized using a regular grid of size h , and our numerical method is based on a standard five-point finite difference scheme. In problems with mixed boundary conditions (Dirichlet and Robin), the Dirichlet boundary condition is enforced weakly using a penalty parameter ε of order 10^{-10} . Unless otherwise stated, the mesh size will be set to $h = 2 \cdot 10^{-2}$ in all experiments. In addition, each convergence analysis is performed taking u_{ex} (exact solution) as the discrete solution obtained when solving on the whole domain Ω with a direct solver.

4.1 Example 1

In order to illustrate the result of Theorem 6, i.e. convergence for the even symmetric part, we take the source term $f = f_e = 1$ in Ω , and set the Dirichlet boundary condition to $g = g_e = 0$ on $\partial\Omega$. A simple initial guess compatible with the Dirichlet boundary condition is in this case $\lambda^0 = \lambda_e^0 = 0$ on Γ . The results are displayed in Figure 3. As expected, when $\theta = \frac{1}{2}$, the DN method becomes a direct solver, thus the error at iteration 2 is "zero" (here it cannot be smaller than the order of magnitude of ε), see Figure 3b. For $\theta \neq \frac{1}{2}$, we see from Figure 3c and Figure 3d that the error on Ω is multiplied by a constant from one iteration to the next. Moreover, the plot of the L^2 and broken H^1 norms of the error in Figure 3e confirms that the DN method converges geometrically in this case. We also see that, as predicted by the proof of Theorem 6, at each iteration, the error remains continuous at the cross-point from Ω_1 to Ω_3 , and from Ω_2 to Ω_4 , and it has the expected symmetry property.

4.2 Example 2

We illustrate now the result of Theorem 13 for the even symmetric part with Robin conditions: the source term is $f = f_e = 1$ in Ω , and the Robin boundary condition is defined by $p = p_e = 1$ and $g = g_e$ on $\partial\Omega$, where g_e is such that $g_e = 1$ on $\partial\Omega_b \cup \partial\Omega_t$ (bottom and top sides) and $g_e = y^2$ on $\partial\Omega_l \cup \partial\Omega_r$ (left and right sides). The same initial guess $\lambda^0 = \lambda_e^0 = 0$ on Γ is considered. As shown in Figure 4, we observe the same convergence properties as for the Dirichlet problem (Example 1). Especially, the DN method becomes a direct




Fig. 3: Results for the DN method applied to (6) (Example 1).

solver when θ is set to $\frac{1}{2}$ (see Figure 4b), and for other choices of θ , it converges geometrically (see Figure 4e), with the expected common ratio $(1 - 2\theta)$.

4.3 Example 3

Finally, we give an illustration of the problematic case described in Theorem 7 and Theorem 8. We consider a Dirichlet problem with the odd symmetric data: $f = f_o = \frac{5\pi^2}{4} \sin(\pi x) \cos(\frac{\pi}{2}y)$ in Ω and $g = g_o = 0$ on $\partial\Omega$. As in




Fig. 4: Results for the DN method applied to the even symmetric part of (2) (Example 2).

the previous examples, the initial guess is set to $\lambda^0 = \lambda_o^0 = 0$. The results displayed in Figure 5 show that, as expected, the DN method applied to this odd symmetric problem does not converge. More specifically, we see that after the first iteration (see Figure 5b), continuity across the cross-point from Ω_2 to Ω_4 is already lost. This jump (referred to as $\delta^1 \neq 0$ in the proof of Theorem 7) generates a "singularity" at the next iteration, see Figure 5c. This "singularity" keeps propagating in the following iterations so that the method diverges, as




Fig. 5: Results for the DN method applied to (7) (Example 3).

predicted by Theorem 8. Moreover, the graph in Figure 5d reveals a geometric (divergent) behaviour of the errors in L^2 and broken H^1 norms.

Remark 4 Since we use a standard finite difference scheme, it is not possible to enforce a discontinuous Dirichlet boundary condition. In practice, when two Dirichlet boundary values do not match at a corner, the average is computed and imposed at this corner. Thus every numerical solution in Ω_i necessarily belongs to a finite dimensional subspace of $C^0(\overline{\Omega_i})$, which explains the quotation marks for singularity in the previous paragraph. More generally, solving this problem using standard discretization methods (such as the finite difference method or the finite element method) involves a regularization step, namely the projection of the discontinuous boundary condition onto some finite dimensional subspace of $C^0(\overline{\Omega_i})$.

Given our choice of projection (computing the average), the boundary data are regularized in the disk D_h of radius h centered at the origin. Outside this disk, they are not modified. Consequently, one may argue that, in each subdomain Ω_i , the local numerical solution should be "not too far" from the local real solution in $\Omega_i \setminus D_h$. In order to verify this in the numerical results, we




Fig. 6: Error at the cross-point at iteration 2 with respect to h , for the DN method applied to the odd symmetric problem described in Example 3, with $\theta = 0.45$.

have plotted (red marks) the value of the error at the cross-point in subdomain Ω_2 with respect to the mesh size h , see Figure 6. We see that these points follow a curve of $(\ln h)^2$ type. An interpretation of that result is that the discretization process (which acts as a regularization here) turns the singularity of type $(\ln r)^2$ from Theorem 7 (see formula (17)) into a pseudo "singularity" of type $(\ln h)^2$.

Now, we would like to analyze how the "singularity" propagates through the numerical iterates. In other words, given the "singularity" of type $(\ln h)^2$ obtained at iteration $k = k_0$, do we observe a "singularity" of type $(\ln h)^{2p+2}$ at the following iterations $k = k_0 + p$ with $p > 0$, as predicted by Theorem 8? To answer this, let us first note that the error at the cross-point $|e_{o,2}^k(0,0)|$ seems to grow geometrically with respect to k , see Figure 5d. Thus, for each h , we are able to compute constants α_2, β_2 such that, for $k \geq 2$,

$$\ln |e_{o,2}^k(0,0)| \simeq \alpha_2 k + \beta_2.$$

In addition, computing the logarithm of formula (25) (with $i = 2$), we get in the neighbourhood of the cross-point

$$\ln |e_{o,2}^k| \simeq \ln \left(\theta \left(\frac{4}{\pi} \ln r \right)^2 \right) k + \tilde{\beta}_2,$$

where $\tilde{\beta}_2$ depends on θ and k (only logarithmically). In Figure 7, we have plotted the value computed for the coefficient α_2 as a function of the mesh size h , and tried to make it fit with a curve of type $\ln(c_1(\ln h)^2)$ (drawn in




Fig. 7: Slope of the curve $k \mapsto \ln |e_{o,2}^k(0,0)|$ with respect to h , for the DN method applied to the odd symmetric problem described in Example 3, with $\theta = 0.45$.

orange). As shown in the figure, this fitting was not successful and it appears that the appropriate fitting is a curve of type $c_2 \ln(c_3(\ln h)^2)$ (drawn in blue), with a constant $c_2 \simeq 0.87$. This suggests that, in the numerical experiments, the "singularity" at iteration $k = k_0 + p$ is of type $(\ln h)^{1.74(p+1)}$ rather than $(\ln h)^{2(p+1)}$. One possible explanation for this is the regularizing effect of the discretization, which may slow down the propagation of the "singularity".

5 Conclusion

We presented a complete analysis of the Dirichlet-Neumann method at the continuous level in a specific configuration involving one cross-point. Based on an even/odd symmetric decomposition of the data, we proved that the even symmetric part of the iterates converges geometrically to the right solution, while the boundary value problems associated to the odd symmetric part are not well-posed, which generates singular iterates. We also exhibited the type of singularity generated, and showed how this singularity propagates through the iterations. Finally, we studied the impact of our theoretical findings on numerical experiments.

A natural extension of this work would be to conduct a similar analysis for the Neumann-Neumann method, which is also known to pose problems in configurations with cross-points, see [41]. Another direction of future work, which will be the subject of the second part of this paper, is to build a modified (and convergent) Dirichlet-Neumann method taking advantage of this even/odd symmetric decomposition of the data.

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

The authors declare that they have no conflict of interest.

References

- [1] Schwarz, H.A.: Über einen Grenzübergang durch alternierendes Verfahren. Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich **15**, 272–286 (1870)
- [2] Bjørstad, P.E., Widlund, O.B.: Iterative methods for the solution of elliptic problems on regions partitioned into substructures. SIAM J. Numer. Anal. **23**(6), 1093–1120 (1986)
- [3] Bourgat, J.-F., Glowinski, R., Le Tallec, P., Vidrascu, M.: Variational formulation and algorithm for trace operator in domain decomposition calculations. In: Chan, T., Glowinski, R., Périaux, J., Widlund, O. (eds.) Domain Decomposition Methods, pp. 3–16. SIAM, Philadelphia, PA (1989)
- [4] Farhat, C., Roux, F.-X.: A method of Finite Element Tearing and Interconnecting and its parallel solution algorithm. Int. J. Numer. Meth. Engrg. **32**, 1205–1227 (1991)
- [5] Toselli, A., Widlund, O.: Domain Decomposition Methods - Algorithms and Theory. Springer Series in Computational Mathematics, vol. 34. Springer, Berlin (2004)
- [6] Hackbusch, W.: Ein iteratives Verfahren zur schnellen Auflösung elliptischer randwertprobleme, rep. 76-12. Technical Report 76-12, Institute for Applied Mathematics, University of Cologne, West Germany, Cologne (1976)
- [7] Brandt, A.: Multi-level adaptive solutions to boundary value problems. Math. Comp. **31**, 333–390 (1977)
- [8] Quarteroni, A., Valli, A.: Domain Decomposition Methods for Partial Differential Equations. Oxford University Press, Oxford (1999)
- [9] Lions, P.-L.: On the Schwarz alternating method. I. In: Glowinski, R., Golub, G.H., Meurant, G.A., Périaux, J. (eds.) First International Symposium on Domain Decomposition Methods for Partial Differential Equations, pp. 1–42. SIAM, Philadelphia, PA (1988)
- [10] Efstathiou, E., Gander, M.J.: Why Restricted Additive Schwarz converges faster than Additive Schwarz. BIT Numerical Mathematics **43**(5), 945–959 (2003)

- [11] Gander, M.J.: Schwarz methods over the course of time. *Electron. Trans. Numer. Anal* **31**(5), 228–255 (2008)
- [12] Després, B.: Méthodes de décomposition de domaine pour les problèmes de propagation d’ondes en régimes harmoniques. PhD thesis, Paris IX (1991)
- [13] Lions, P.-L.: On the Schwarz alternating method. III: a variant for nonoverlapping subdomains. In: Chan, T.F., Glowinski, R., Périaux, J., Widlund, O. (eds.) *Third International Symposium on Domain Decomposition Methods for Partial Differential Equations*, Held in Houston, Texas, March 20–22, 1989. SIAM, Philadelphia, PA (1990)
- [14] Boubendir, Y., Bendali, A.: Dealing with cross-points in a non-overlapping domain decomposition solution of the Helmholtz equation. In: *Mathematical and Numerical Aspects of Wave Propagation WAVES 2003*, pp. 319–324. Springer, Jyväskylä (2003)
- [15] Gander, M.J., Kwok, F.: Best Robin parameters for optimized Schwarz methods at cross points. *SIAM Journal on Scientific Computing* **34**(4), 1849–1879 (2012)
- [16] Loisel, S.: Condition number estimates for the nonoverlapping optimized Schwarz method and the 2-Lagrange multiplier method for general domains and cross points. *SIAM Journal on Numerical Analysis* **51**(6), 3062–3083 (2013)
- [17] Gander, M.J., Kwok, F.: On the applicability of Lions’ energy estimates in the analysis of discrete optimized Schwarz methods with cross points. In: *Domain Decomposition Methods in Science and Engineering XX*, pp. 475–483. Springer, San Diego, CA (2013)
- [18] Gander, M.J., Santugini-Repiquet, K.: Cross-points in domain decomposition methods with a finite element discretization. *ETNA* **45**, 219–240 (2016)
- [19] Bamberger, A., Joly, P., Roberts, J.E.: Second-order absorbing boundary conditions for the wave equation: a solution for the corner problem. *SIAM J. Num. Anal.* **27**(2), 323–352 (1990)
- [20] Nataf, F.: A Schwarz additive method with high order interface conditions and nonoverlapping subdomains. *ESAIM: Mathematical Modelling and Numerical Analysis-Modélisation Mathématique et Analyse Numérique* **32**(1), 107–116 (1998)
- [21] Modave, A., Royer, A., Antoine, X., Geuzaine, C.: A non-overlapping domain decomposition method with high-order transmission conditions

- and cross-point treatment for Helmholtz problems. *Computer Methods in Applied Mechanics and Engineering* **368**, 113162 (2020)
- [22] Després, B., Nicolopoulos, A., Thierry, B.: Corners and stable optimized domain decomposition methods for the Helmholtz problem. hal-02612368 (2020)
 - [23] Gander, M.J., Halpern, L.: A simple finite difference discretization for Ventcell transmission conditions at cross points. In: *Domain Decomposition Methods in Science and Engineering XXVI*. Springer, Hong-Kong (2021)
 - [24] Chaouqui, F., Gander, M.J., Santugini-Repiquet, K.: A local coarse space correction leading to a well-posed continuous neumann-neumann method in the presence of cross points. In: *International Conference on Domain Decomposition Methods*, pp. 83–91 (2018). Springer
 - [25] Claeys, X., Parolin, E.: Robust treatment of cross points in optimized Schwarz methods. arXiv preprint arXiv:2003.06657 (2020)
 - [26] Gander, M.J., Kwok, F.: Optimal interface conditions for an arbitrary decomposition into subdomains. In: *Domain Decomposition Methods in Science and Engineering XIX*, pp. 101–108. Springer, Zhangjiajie (2011)
 - [27] *Domain Decomposition Methods in Science and Engineering XXVI*. Springer, Hong-Kong (2021)
 - [28] De Roeck, Y.-H., Le Tallec, P.: Analysis and test of a local domain-decomposition preconditioner. In: *Domain Decomposition Methods in Science and Engineering IV*. Springer, Moscow (1990)
 - [29] Maier, I., Haasdonk, B.: A dirichlet–neumann reduced basis method for homogeneous domain decomposition problems. *Applied Numerical Mathematics* **78**, 31–48 (2014)
 - [30] Song, B., Jiang, Y.-L., Wang, X.: Analysis of two new parareal algorithms based on the dirichlet-neumann/neumann-neumann waveform relaxation method for the heat equation. *Numerical Algorithms* **86**(4), 1685–1703 (2021)
 - [31] Chaouqui, F., Ciaramella, G., Gander, M.J., Vanzan, T.: On the scalability of classical one-level domain-decomposition methods. *Vietnam Journal of Mathematics* **46**(4), 1053–1088 (2018)
 - [32] Gander, M.J., Kwok, F., Mandal, B.C.: Dirichlet–neumann waveform relaxation methods for parabolic and hyperbolic problems in multiple subdomains. *BIT Numerical Mathematics* **61**(1), 173–207 (2021)

- [33] Widlund, O., Dryja, M., Proskurowski, W.: A method of domain decomposition with crosspoints for elliptic finite element problems. In: Proceedings of the International Symposium on Optimal Algorithms. Blagovgrad, Bulgaria, April 21-25, 1986, pp. 97–111 (1986). Bulgarian Academy of Sciences, Sofia
- [34] Dryja, M.: A method of domain decomposition for three-dimensional finite element elliptic problems. In: First International Symposium on Domain Decomposition Methods for Partial Differential Equations, pp. 43–61 (1988). SIAM Philadelphia
- [35] Parolin, É.: Non-overlapping domain decomposition methods with non-local transmission operators for harmonic wave propagation problems. PhD thesis, Institut Polytechnique de Paris (2020)
- [36] Dauge, M.: Elliptic Boundary Value Problems on Corner Domains: Smoothness and Asymptotics of Solutions vol. 1341. Springer, Berlin (2006)
- [37] Grisvard, P.: Elliptic Problems in Nonsmooth Domains. SIAM, Philadelphia, PA (2011)
- [38] Dauge, M.: Regularity and singularities in polyhedral domains. Available on: https://perso.univ-rennes1.fr/monique.dauge/publis/Talk_Karlsruhe08.pdf (2008)
- [39] Demlow, A.: Lecutre notes: Singular solutions to pde and their finite element approximations. Available on: https://www.math.tamu.edu/~demlow/Courses/16_sp/663_sp16/index.html (2016)
- [40] Brézis, H.: Functional Analysis, Sobolev Spaces and Partial Differential Equations vol. 2. Springer, New-York (2011)
- [41] Chaouqui, F., Gander, M.J., Santugini-Repiquet, K.: A coarse space to remove the logarithmic dependency in neumann-neumann methods. In: International Conference on Domain Decomposition Methods, pp. 159–167 (2017). Springer