



UNIVERSITÉ
DE GENÈVE

PRESS RELEASE

Geneva | November 16th, 2020



Hôpitaux
Universitaires
Genève



UNIVERSITÀ DEGLI STUDI DI NAPOLI
FEDERICO II



FATEBENEFRATELLI

SYNLAB
IRCCS SDN

Link between Alzheimer's disease and gut microbiota is confirmed

Swiss and Italian scientists prove a correlation between gut microbiota and the appearance of amyloid plaques in the brain, typical of Alzheimer's disease.

UNIVERSITÉ DE GENÈVE
Communication Department

24 rue du Général-Dufour
CH-1211 Geneva 4


Tel. +41 22 379 77 17
media@unige.ch
www.unige.ch

Alzheimer's disease is the most common cause of dementia. Still incurable, it directly affects nearly one million people in Europe, and indirectly millions of family members as well as society as a whole. In recent years, the scientific community has suspected that the gut microbiota plays a role in the development of the disease. A team from the University of Geneva (UNIGE) and the University Hospitals of Geneva (HUG) in Switzerland, together with Italian colleagues from the National Research and Care Center for Alzheimer's and Psychiatric Diseases Fatebenefratelli in Brescia, University of Naples and the IRCCS SDN Research Center in Naples, confirm the correlation, in humans, between an imbalance in the gut microbiota and the development of amyloid plaques in the brain, which are at the origin of the neurodegenerative disorders characteristic of Alzheimer's disease. Proteins produced by certain intestinal bacteria, identified in the blood of patients, could indeed modify the interaction between the immune and the nervous systems and trigger the disease. These results, to be discovered in the *Journal of Alzheimer's Disease*, make it possible to envisage new preventive strategies based on the modulation of the microbiota of people at risk.

The research laboratory of neurologist Giovanni Frisoni, director of the HUG Memory Centre and professor at the Department of Rehabilitation and Geriatrics of the UNIGE Faculty of Medicine, has been working for several years now on the potential influence of the gut microbiota on the brain, and more particularly on neurodegenerative diseases. «We have already shown that the gut microbiota composition in patients with Alzheimer's disease was altered, compared to people who do not suffer from such disorders,» he explains. «Their microbiota has indeed a reduced microbial diversity, with an over-representation of certain bacteria and a strong decrease in other microbes. Furthermore, we have also discovered an association between an inflammatory phenomenon detected in the blood, certain intestinal bacteria and Alzheimer's disease; hence the hypothesis that we wanted to test here: could inflammation in the blood be a mediator between the microbiota and the brain? »

The brain under influence

Intestinal bacteria can influence the functioning of the brain and promote neurodegeneration through several pathways: they can indeed influence the regulation of the immune system and, consequently, can modify the interaction between the immune system and the nervous system. Lipopolysaccharides, a protein located on the membrane of bacteria with pro-inflammatory properties, have been found



The gut microbiota could play a role in brain diseases including Alzheimer's disease. Some gut bacteria release lipopolysaccharides and short chain fatty acids that can reach the brain via the blood and might cause amyloid pathology and neurodegeneration.

High resolution pictures

contact

Giovanni Frisoni

Full Professor
Department of Rehabilitation
and Geriatrics
Faculty of Medicine, UNIGE
Director of the Memory Centre,
HUG

+41 22 372 58 00

Giovanni.Frisoni@unige.ch

Moira Marizzoni

Laboratory of Biological Psychiatry and Laboratory of Neuroimaging and Alzheimer's Epidemiology, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy

+39 030 35 01 362

mmarizzoni@fatebenefratelli.eu

DOI: [10.3233/JAD-200306](https://doi.org/10.3233/JAD-200306)

in amyloid plaques and around vessels in the brains of people with Alzheimer's disease. In addition, the intestinal microbiota produces metabolites - in particular some short-chain fatty acids - which, having neuroprotective and anti-inflammatory properties, directly or indirectly affect brain function.

«To determine whether inflammation mediators and bacterial metabolites constitute a link between the gut microbiota and amyloid pathology in Alzheimer's disease, we studied a cohort of 89 people between 65 and 85 years of age. Some suffered from Alzheimer's disease or other neurodegenerative diseases causing similar memory problems, while others did not have any memory problems,» reports Moira Marizzoni, a researcher at the Fatebenefratelli Center in Brescia and first author of this work. «Using PET imaging, we measured their amyloid deposition and then quantified the presence in their blood of various inflammation markers and proteins produced by intestinal bacteria, such as lipopolysaccharides and short-chain fatty acids.»

A very clear correlation

«Our results are indisputable: certain bacterial products of the intestinal microbiota are correlated with the quantity of amyloid plaques in the brain,» explains Moira Marizzoni. «Indeed, high blood levels of lipopolysaccharides and certain short-chain fatty acids (acetate and valerate) were associated with both large amyloid deposits in the brain. Conversely, high levels of another short-chain fatty acid, butyrate, were associated with less amyloid pathology.»

This work thus provides proof of an association between certain proteins of the gut microbiota and cerebral amyloidosis through a blood inflammatory phenomenon. Scientists will now work to identify specific bacteria, or a group of bacteria, involved in this phenomenon.

A strategy based on prevention

This discovery paves the way for potentially highly innovative protective strategies - through the administration of a bacterial cocktail, for example, or of pre-biotics to feed the «good» bacteria in our intestine. «However, we shouldn't be too quick to rejoice,» says Frisoni. «Indeed, we must first identify the strains of the cocktail. Then, a neuroprotective effect could only be effective at a very early stage of the disease, with a view to prevention rather than therapy. However, early diagnosis is still one of the main challenges in the management of neurodegenerative diseases, as protocols must be developed to identify high-risk individuals and treat them well before the appearance of detectable symptoms.» This study is also part of a broader prevention effort led by the UNIGE Faculty of Medicine and the HUG Memory Centre.

This study was supported by the Swiss Association for Alzheimer's Research, the Segre Foundation, the Velux Foundation, the Edmond J. Safra Foundation, the Fondazione Agusta, the Chmielewski Foundation, Mr. Ivan Pictet, as well as other funding institutions.