

Noninteracting fermions in a 2d rotating trap and random matrix theory

Grégory Schehr, LPTHE
CNRS/Sorbonne Université, Paris

*Random matrices and random landscapes,
24-29 July 2022*

Noninteracting fermions in a 2d rotating trap and random matrix theory

Grégory Schehr, LPTHE
CNRS/Sorbonne Université, Paris

*Random matrices and random landscapes,
24-29 July 2022*

- Manas Kulkarni (ICTS-TIFR, Bangalore)
- Bertrand Lacroix-A-Chez-Toine (King 's College, London)
- Pierre Le Doussal (LPENS, Ecole Normale Sup., Paris)
- Satya N. Majumdar (LPTMS, Univ. Paris-Saclay)
- Naftali R. Smith (Ben-Gurion Univ., Israel)

Noninteracting fermions in a 2d rotating trap and random matrix theory

Grégory Schehr, LPTHE
CNRS/Sorbonne Université, Paris

*Random matrices and random landscapes,
24-29 July 2022*

- Manas Kulkarni (ICTS-TIFR, Bangalore)
- Bertrand Lacroix-A-Chez-Toine (King 's College, London)
- Pierre Le Doussal (LPENS, Ecole Normale Sup., Paris)
- Satya N. Majumdar (LPTMS, Univ. Paris-Saclay)
- Naftali R. Smith (Ben-Gurion Univ., Israel)


Lacroix-A-Chez-Toine, Majumdar, G. S., PRA **99**, 021602 (R), (2019)

Kulkarni, Majumdar, G. S., PRA **103**, 033321 (2021)

Smith, Le Doussal, Majumdar, G. S., PRA **105**, 043315 (2022)

Ultra-cold atoms in confining potentials

- Recent progress in the experimental manipulation of cold atoms




to investigate the interplay between **quantum** and **thermal** behaviors in many-body systems at low temperature

Ultra-cold atoms in confining potentials


- Recent progress in the experimental manipulation of cold atoms

➔ to investigate the interplay between **quantum** and **thermal** behaviors in many-body systems at low temperature

- A common feature of these experiments: presence of a **confining potential** that traps the atoms within a limited spatial region




Ultra-cold atoms in confining potentials




- **bulk**: traditional many-body physics (**translationally** invariant system)

Ultra-cold atoms in confining potentials



- **bulk**: traditional many-body physics (**translationally** invariant system)


Ultra-cold atoms in confining potentials



- **bulk**: traditional many-body physics (**translationally** invariant system)
«The uniform electron gas, the traditional starting point for density-based many-body theories of inhomogeneous systems, is inappropriate near electronic edges.»

W. Kohn, A. E. Mattsson, PRL 1998


Ultra-cold atoms in confining potentials



- **bulk**: traditional many-body physics (**translationally** invariant system)

This talk: random matrix theory is the ideal tool to study these edge properties


Ultra-cold atoms in confining potentials



- **bulk**: traditional many-body physics (**translationally** invariant system)
- **edge**: new physics induced by **confinement** → **universal edge** properties


This talk: random matrix theory is the ideal tool to study these edge properties

Spinless free fermions in a 1d harmonic potential



$$V(x) = \frac{1}{2}m\omega^2 x^2$$

Spinless free fermions in a 1d harmonic potential




$$V(x) = \frac{1}{2}m\omega^2 x^2$$

At zero temperature: connection between spinless free fermions in a harmonic trap and Random Matrix Theory (GUE)

Connection between free fermions at $T=0$ and RMT

- A single quantum particle in a harmonic potential




$$V(x) = \frac{1}{2}m\omega^2 x^2$$

$$\hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + \frac{1}{2}m\omega^2 x^2$$

Connection between free fermions at T=0 and RMT

- A single quantum particle in a harmonic potential



$$V(x) = \frac{1}{2}m\omega^2 x^2$$

$$\hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + \frac{1}{2}m\omega^2 x^2$$

- Single particle eigenfunctions

$$\hat{H} \varphi_E(x) = E \varphi_E(x)$$

with $\varphi_E(x \rightarrow \pm\infty) = 0$


$$\varphi_k(x) = \left[\frac{\alpha}{\sqrt{\pi} 2^k k!} \right]^{1/2} e^{-\frac{\alpha^2 x^2}{2}} H_k(\alpha x)$$

$$\epsilon_k = \hbar\omega(k + 1/2) \quad , \quad \alpha = \sqrt{m\omega/\hbar}$$

$$k \in \mathbb{N}$$

Connection between free fermions at T=0 and RMT

- A single quantum particle in a harmonic potential



$$V(x) = \frac{1}{2}m\omega^2 x^2$$

$$\hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + \frac{1}{2}m\omega^2 x^2$$

Hermite polynomial

- Single particle eigenfunctions

$$\hat{H} \varphi_E(x) = E \varphi_E(x)$$


with $\varphi_E(x \rightarrow \pm\infty) = 0$

$$\varphi_k(x) = \left[\frac{\alpha}{\sqrt{\pi} 2^k k!} \right]^{1/2} e^{-\frac{\alpha^2 x^2}{2}} H_k(\alpha x)$$


$$\epsilon_k = \hbar\omega(k + 1/2) \quad , \quad \alpha = \sqrt{m\omega/\hbar}$$

$$k \in \mathbb{N}$$


N spinless free fermions in a 1d harmonic trap at $T=0$




N spinless free fermions in a 1d harmonic trap at $T=0$




N spinless free fermions in a 1d harmonic trap at $T=0$




N spinless free fermions in a 1d harmonic trap at $T=0$




N spinless free fermions in a 1d harmonic trap at $T=0$




N spinless free fermions in a 1d harmonic trap at $T=0$



N spinless free fermions in a 1d harmonic trap at $T=0$



N spinless free fermions in a 1d harmonic trap at $T=0$




- The N -particle wave function is given by a $N \times N$ **Slater determinant**

$$\Psi_0(x_1, x_2, \dots, x_N) = \frac{1}{\sqrt{N!}} \det[\varphi_i(x_j)] \quad \begin{matrix} 0 \leq i \leq N-1 \\ 1 \leq j \leq N \end{matrix}$$

$$\varphi_k(x) = \left[\frac{\alpha}{\sqrt{\pi} 2^k k!} \right]^{1/2} e^{-\frac{\alpha^2 x^2}{2}} H_k(\alpha x)$$

N spinless free fermions in a 1d harmonic trap at $T=0$




- The N -particle wave function is given by a $N \times N$ **Slater determinant**

$$\Psi_0(x_1, x_2, \dots, x_N) = \frac{1}{\sqrt{N!}} \det[\varphi_i(x_j)] \quad \begin{matrix} 0 \leq i \leq N-1 \\ 1 \leq j \leq N \end{matrix}$$

$$\varphi_k(x) = \left[\frac{\alpha}{\sqrt{\pi} 2^k k!} \right]^{1/2} e^{-\frac{\alpha^2 x^2}{2}} H_k(\alpha x)$$


$$\Rightarrow \Psi_0(x_1, x_2, \dots, x_N) \propto e^{-\frac{\alpha^2}{2}(x_1^2 + \dots + x_N^2)} \prod_{1 \leq j < k \leq N} (x_j - x_k)$$

N spinless free fermions in a 1d harmonic trap at $T=0$



- Probability density function (PDF) of the positions

$$|\Psi_0(x_1, \dots, x_N)|^2 = \frac{1}{z_N(\alpha)} \prod_{i < j} (x_i - x_j)^2 e^{-\alpha^2 \sum_{i=1}^N x_i^2}$$



The positions of the free fermions behave statistically like the eigenvalues of GUE random matrices


Properties of fermions in a 1d harmonic trap at $T=0$

- Average density of free fermions: **Wigner semi-circle law**

$$\rho_N(x, T=0) = \frac{1}{N} \sum_{i=1}^N \langle \delta(x - x_i) \rangle$$

for $N \gg 1$ $\rho_N(x, T=0) \approx \frac{\alpha}{\sqrt{N}} f_W\left(\frac{\alpha x}{\sqrt{N}}\right)$, $f_W(z) = \frac{1}{\pi} \sqrt{2 - z^2}$

See also Local Density (or Thomas-Fermi) Approx. in the literature on fermions



What about other Hermitian matrix models ?

- **Laguerre Unitary Ensemble** can be realized with a 1d singular potential

$$V(x) = \frac{\alpha(\alpha - 1)}{x^2} + \beta x^2, \quad x > 0$$

Nadal, Majumdar, PRE '09

- **Jacobi Unitary Ensemble** can be realized with a 1d box potential

$$V(x) = \begin{cases} 0, & -1 \leq x \leq +1 \\ +\infty, & |x| > 1 \end{cases}$$

Lacroix-A-Chez-Toine, Le Doussal, Majumdar, G. S., EPL '17

- For a review see Dean, Le Doussal, Majumdar, G. S., J. Phys. A '19

What about other Hermitian matrix models ?

- **Laguerre Unitary Ensemble** can be realized with a 1d singular potential

$$V(x) = \frac{\alpha(\alpha - 1)}{x^2} + \beta x^2, \quad x > 0$$


Nadal, Majumdar, PRE '09

- **Jacobi Unitary Ensemble** can be realized with a 1d box potential

$$V(x) = \begin{cases} 0, & -1 \leq x \leq +1 \\ +\infty, & |x| > 1 \end{cases}$$

Lacroix-A-Chez-Toine, Le Doussal, Majumdar, G. S., EPL '17

- For a review see Dean, Le Doussal, Majumdar, G. S., J. Phys. A '19




What about 2d (non Hermitian) matrix ensembles,
e.g., Ginibre ensemble ?

A natural candidate: fermions in a 2d harmonic trap

- Single particle Hamiltonian

$$\hat{H} = -\frac{\hbar^2}{2m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} m \omega^2 \underbrace{(x^2 + y^2)}_{r^2}$$



A natural candidate: fermions in a 2d harmonic trap



- Single particle Hamiltonian

$$\hat{H} = -\frac{\hbar^2}{2m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} m \omega^2 \underbrace{(x^2 + y^2)}_{r^2}$$

- Global density (at T=0)

$$\rho_N(r) \approx \frac{1}{N} \frac{m}{2\pi\hbar^2} \left(\mu - \frac{1}{2} m \omega^2 r^2 \right) \quad \text{with} \quad \mu \approx \hbar\omega\sqrt{2N}$$

spatial extent $R_e \propto N^{1/4}$



Dean, Le Doussal, Majumdar, G. S., EPL '15, PRA '16
see also G. Lambert's talk

A natural candidate: fermions in a 2d harmonic trap



- Single particle Hamiltonian

$$\hat{H} = -\frac{\hbar^2}{2m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} m \omega^2 \underbrace{(x^2 + y^2)}_{r^2}$$

- Global density (at T=0)

$$\rho_N(r) \approx \frac{1}{N} \frac{m}{2\pi\hbar^2} \left(\mu - \frac{1}{2} m \omega^2 r^2 \right) \quad \text{with} \quad \mu \approx \hbar\omega\sqrt{2N}$$

spatial extent $R_e \propto N^{1/4}$



Dean, Le Doussal, Majumdar, G. S., EPL '15, PRA '16
see also G. Lambert's talk

No apparent connection with known
matrix models !


Fermions in a 2D rotating harmonic trap

- Single particle Hamiltonian in the rotating frame

$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$

where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum

and $L_z = i(y\partial_x - x\partial_y)$



Fermions in a 2D rotating harmonic trap


- Single particle Hamiltonian in the rotating frame

$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$

where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum

and $L_z = i(y\partial_x - x\partial_y)$

$$\hat{H} = \frac{\mathbf{p}^2}{2} + \frac{1}{2} \omega^2 \mathbf{r}^2 - \Omega L_z$$



Fermions in a 2D rotating harmonic trap

- Single particle Hamiltonian in the rotating frame

$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$


where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum

and $L_z = i(y\partial_x - x\partial_y)$

$$\hat{H} = \frac{\mathbf{p}^2}{2} + \frac{1}{2} \omega^2 \mathbf{r}^2 - \Omega L_z$$

- Analogy with a magnetic system

$$\hat{H} = \frac{1}{2} (\mathbf{p} - \mathbf{A})^2 + \frac{1}{2} \omega^2 \mathbf{r}^2 - \frac{1}{2} \Omega^2 \mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap

- Single particle Hamiltonian in the rotating frame


$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$

where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum

and $L_z = i(y\partial_x - x\partial_y)$

$$\hat{H} = \frac{\mathbf{p}^2}{2} + \frac{1}{2} \omega^2 \mathbf{r}^2 - \Omega L_z$$

- Analogy with a magnetic system



Coriolis
potential

$$\hat{H} = \frac{1}{2} (\mathbf{p} - \mathbf{A})^2 + \frac{1}{2} \omega^2 \mathbf{r}^2 - \frac{1}{2} \Omega^2 \mathbf{r}^2$$

Fermions in a 2D rotating harmonic trap

- Single particle Hamiltonian in the rotating frame


$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$

where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum

and $L_z = i(y\partial_x - x\partial_y)$

$$\hat{H} = \frac{\mathbf{p}^2}{2} + \frac{1}{2} \omega^2 \mathbf{r}^2 - \Omega L_z$$

- Analogy with a magnetic system



Coriolis
potential

centrifugal
potential

$$\hat{H} = \frac{1}{2} (\mathbf{p} - \mathbf{A})^2 + \frac{1}{2} \omega^2 \mathbf{r}^2 - \frac{1}{2} \Omega^2 \mathbf{r}^2$$

Fermions in a 2D rotating harmonic trap

- Single particle Hamiltonian in the rotating frame

$$\hat{H} = -\frac{1}{2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + \frac{1}{2} \omega^2 (x^2 + y^2) - \Omega L_z$$

where $\mathbf{L} = (\mathbf{r} \wedge \mathbf{p})$: angular momentum


and $L_z = i(y\partial_x - x\partial_y)$

$$\hat{H} = \frac{\mathbf{p}^2}{2} + \frac{1}{2} \omega^2 \mathbf{r}^2 - \Omega L_z$$

- Analogy with a magnetic system

effective mag. field

$$\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$$



Coriolis potential

centrifugal potential

$$\hat{H} = \frac{1}{2} (\mathbf{p} - \mathbf{A})^2 + \frac{1}{2} \omega^2 \mathbf{r}^2 - \frac{1}{2} \Omega^2 \mathbf{r}^2$$


$$\mathbf{A} = \begin{pmatrix} \Omega y \\ -\Omega x \end{pmatrix}$$

symmetric gauge

Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$


$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$



$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$




$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$



$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$


$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$





Fermions in a 2D rotating harmonic trap

- Analogy with a magnetic system $\mathbf{B}_{\text{eff}} = \nabla \wedge \mathbf{A} = \Omega \mathbf{u}_z$

$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



the Fermi gas becomes unstable



Fermions in a 2D rotating harmonic trap

$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\mathbf{r}^2 - \frac{1}{2}\Omega^2\mathbf{r}^2 \quad \text{we set} \quad \hbar = m = \omega = 1$$


- Single particle eigenfunctions $\psi_{k,\ell}(r, \theta)$ in polar coordinates are labelled by two quantum numbers (k, ℓ) with $k \in \mathbb{N}$ and $\ell \in \mathbb{Z}$

$$\hat{H}\psi_{k,\ell} = \epsilon_{k,\ell}\psi_{k,\ell} \quad \left\{ \begin{array}{l} \psi_{k,\ell}(r, \theta) = a_{k,\ell} L_k^{|\ell|}(r^2) r^{|\ell|} e^{-r^2/2} e^{i\ell\theta} \\ E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell \end{array} \right.$$

Fermions in a 2D rotating harmonic trap: Landau levels

$$\hbar = m = \omega = 1$$


$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: Landau levels

$$\hbar = m = \omega = 1$$


$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: Landau levels

$$\hbar = m = \omega = 1$$


$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: Landau levels

$$\hbar = m = \omega = 1$$


$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: Landau levels

$$\hbar = m = \omega = 1$$


$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: Landau levels


$$\hbar = m = \omega = 1$$

$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$



Fermions in a 2D rotating harmonic trap: ground state


- Ground state of N noninteracting spinless fermions



- If $E_{0,\ell=N-1} < E_{0,\ell=-1} \simeq E_{1,\ell=0}$ only the **LLL** is occupied

Fermions in a 2D rotating harmonic trap: ground state


- Ground state of N noninteracting spinless fermions



- If $E_{0,\ell=N-1} < E_{0,\ell=-1} \simeq E_{1,\ell=0}$ only the LLL is occupied

Fermions in a 2D **rotating** harmonic trap: ground state


- Ground state of N noninteracting spinless fermions



- Only the **LLL** is occupied $\iff \omega \left(1 - \frac{2}{N}\right) < \Omega < \omega$

Fermions in a 2D rotating harmonic trap: ground state

- Ground state of N noninteracting spinless fermions



- Only the LLL is occupied

$$\iff \omega \left(1 - \frac{2}{N} \right) < \Omega < \omega$$

Fermions in a 2D rotating harmonic trap: ground state

- Suppose that only the LLL is occupied $\iff \omega \left(1 - \frac{2}{N}\right) < \Omega < \omega$

Fermions in a 2D rotating harmonic trap: ground state

- Suppose that only the LLL is occupied $\iff \omega \left(1 - \frac{2}{N}\right) < \Omega < \omega$

- In the LLL, the single particle eigenstates/eigenvalues are

$$\psi_\ell(r, \theta) \equiv \psi_{0,\ell}(r, \theta) = a_{0,\ell} r^\ell e^{i\ell\theta} e^{-r^2/2}, \quad \ell \geq 0$$

$$\psi_\ell(z) = \frac{1}{\sqrt{\pi\ell!}} z^\ell e^{-\frac{1}{2}|z|^2} \quad \text{and} \quad E_\ell = 1 + (1 - \Omega)\ell$$

Fermions in a 2D rotating harmonic trap: ground state

- Suppose that only the LLL is occupied $\iff \omega \left(1 - \frac{2}{N}\right) < \Omega < \omega$

- In the LLL, the single particle eigenstates/eigenvalues are

$$\psi_\ell(r, \theta) \equiv \psi_{0,\ell}(r, \theta) = a_{0,\ell} r^\ell e^{i\ell\theta} e^{-r^2/2}, \quad \ell \geq 0$$

$$\psi_\ell(z) = \frac{1}{\sqrt{\pi\ell!}} z^\ell e^{-\frac{1}{2}|z|^2} \quad \text{and} \quad E_\ell = 1 + (1 - \Omega)\ell$$

- Ground state many-body eigenfunction: Slater determinant

$$\begin{aligned} \Psi_0(z_1, z_2, \dots, z_N) &= \frac{1}{\sqrt{N!}} \det_{1 \leq i, j \leq N} \psi_{i-1}(z_j) \\ &\propto e^{-\frac{1}{2} \sum_{j=1}^N |z_j|^2} \det_{1 \leq i, j \leq N} z_j^{i-1} \propto e^{-\frac{1}{2} \sum_{j=1}^N |z_j|^2} \prod_{1 \leq i < j \leq N} (z_i - z_j) \end{aligned}$$

Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19

- Joint proba. density function of the positions of the fermions


$$P_{\text{joint}}(z_1, z_2, \dots, z_N) = |\Psi_0(z_1, z_2, \dots, z_N)|^2 = A_N \prod_{1 \leq i, j \leq N} |z_i - z_j|^2 e^{-\sum_{i=1}^N |z_i|^2}$$

Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19

- Joint proba. density function of the positions of the fermions

$$P_{\text{joint}}(z_1, z_2, \dots, z_N) = |\Psi_0(z_1, z_2, \dots, z_N)|^2 = A_N \prod_{1 \leq i, j \leq N} |z_i - z_j|^2 e^{-\sum_{i=1}^N |z_i|^2}$$



the positions of the fermions are in one to one correspondence with the eigenvalues of a complex Ginibre random matrix

Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19

- Joint proba. density function of the positions of the fermions

$$P_{\text{joint}}(z_1, z_2, \dots, z_N) = |\Psi_0(z_1, z_2, \dots, z_N)|^2 = A_N \prod_{1 \leq i, j \leq N} |z_i - z_j|^2 e^{-\sum_{i=1}^N |z_i|^2}$$

→ the positions of the fermions are in one to one correspondence with the eigenvalues of a complex Ginibre random matrix

→ this provides a simple physical realisation of the Ginibre ensemble of RMT

Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19

- Joint proba. density function of the positions of the fermions

$$P_{\text{joint}}(z_1, z_2, \dots, z_N) = |\Psi_0(z_1, z_2, \dots, z_N)|^2 = A_N \prod_{1 \leq i, j \leq N} |z_i - z_j|^2 e^{-\sum_{i=1}^N |z_i|^2}$$

→ the positions of the fermions are in one to one correspondence with the eigenvalues of a complex Ginibre random matrix

→ this provides a simple physical realisation of the Ginibre ensemble of RMT

→ this connection has been exploited to compute various observables, like the **full counting statistics** or the **Rényi entanglement entropy**

Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19


- Joint proba. density function of the positions of the fermions

$$P_{\text{joint}}(z_1, z_2, \dots, z_N) = |\Psi_0(z_1, z_2, \dots, z_N)|^2 = A_N \prod_{1 \leq i, j \leq N} |z_i - z_j|^2 e^{-\sum_{i=1}^N |z_i|^2}$$

- Large N limit of the density: **Girko's law**

$$\rho_N(z) = \sum_{i=1}^N \delta(z - z_i) \\ \xrightarrow{N \rightarrow \infty} \frac{1}{\pi} \Theta(\sqrt{N} - |z|)$$



spatial extent $R_e \propto N^{1/2}$



Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19



■ Edge density profile



Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19



■ Edge density profile




Connection to the complex Ginibre ensemble GinUE

Lacroix-A-Chez-Toine, Majumdar, G. S., PRA (R) '19

■ Edge density profile




$\rho^{\text{edge}}(u)$



$$\rho^{\text{edge}}(u) = \frac{1}{\pi^{3/2}} \int_{-u}^{\infty} e^{-x^2} dx = \frac{1}{2\pi} \text{erfc}(-u)$$


Fermions in a 2D rotating harmonic trap

$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2\mathbf{r}^2 - \frac{1}{2}\Omega^2\mathbf{r}^2$$




Fermions in a 2D **rotating** harmonic trap

$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2\mathbf{r}^2 - \frac{1}{2}\Omega^2\mathbf{r}^2$$



Fermions in a 2D rotating harmonic trap


$$\hat{H} = \frac{1}{2}(\mathbf{p} - \mathbf{A})^2 + \frac{1}{2}\omega^2 \mathbf{r}^2 - \frac{1}{2}\Omega^2 \mathbf{r}^2$$



What happens when one decreases Ω ?

Decreasing the rotating frequency


- Ground state of N noninteracting spinless fermions



$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$

Decreasing the rotating frequency

- Ground state of N noninteracting spinless fermions



$$E_{k,\ell} = 2k + 1 + |\ell| - \Omega\ell$$


Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21




Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21





Decreasing the rotating frequency: wedding cake structure

Kulkarni, Majumdar, G. S. '21





Decreasing the rotating frequency: higher Landau levels



- Different layers correspond to different/higher Landau levels

Decreasing the rotating frequency: higher Landau levels





- Different layers correspond to different/higher Landau levels
- Edge density of the k -th step/Landau level

Kulkarni, Majumdar, G. S. '21
Smith, Le Doussal, Majumdar, G. S. '22

$$\rho_k^{\text{edge}}(u) = \frac{2^{-k}}{\pi^{3/2} k!} \int_{-u}^{\infty} e^{-x^2} [H_k(x)]^2 dx$$

Decreasing the rotating frequency: higher Landau levels



- Different layers correspond to different/higher Landau levels
- Edge density of the k -th step/Landau level



Kulkarni, Majumdar, G. S. '21
Smith, Le Doussal, Majumdar, G. S. '22

$$\rho_k^{\text{edge}}(u) = \frac{2^{-k}}{\pi^{3/2} k!} \int_{-u}^{\infty} e^{-x^2} [H_k(x)]^2 dx$$

➡ it exhibits k kinks at the k zeros of the k -th Hermite polynomials

see also Dunne '94

Decreasing the rotating frequency: higher Landau levels



- Different layers correspond to different/higher Landau levels
- Edge density of the k -th step/Landau level

Kulkarni, Majumdar, G. S. '21
Smith, Le Doussal, Majumdar, G. S. '22

$$\rho_k^{\text{edge}}(u) = \frac{2^{-k}}{\pi^{3/2} k!} \int_{-u}^{\infty} e^{-x^2} [H_k(x)]^2 dx$$

➔ it exhibits k kinks at the k zeros of the k -th Hermite polynomials

see also Dunne '94


- Connection to Polyanalytic Ginibre Ensemble

see also Haimi, Hedenmalm '13
Fenzel, Lambert '20

Edge scaling limit for **very high** Landau levels

- Edge density profile of the k -th step/Landau level for $k \rightarrow \infty$

Kulkarni, Majumdar, G. S. '21




$$\lim_{k \rightarrow \infty} \rho_k^{\text{edge}}(u = \sqrt{2k}y) = \frac{1}{\pi^2} \left(\frac{\pi}{2} + \sin^{-1}(y) \right)$$

for $-1 < y < 1$

Edge scaling limit for **very high** Landau levels

- Edge density profile of the k -th step/Landau level for $k \rightarrow \infty$

Kulkarni, Majumdar, G. S. '21



$$\lim_{k \rightarrow \infty} \rho_k^{\text{edge}}(u = \sqrt{2k} y) = \frac{1}{\pi^2} \left(\frac{\pi}{2} + \sin^{-1}(y) \right)$$

for $-1 < y < 1$

- **Airy universality class** at the edge of the edge

$$\rho_k^{\text{edge}} \left(-\sqrt{2k} + \frac{w}{\sqrt{2k^{1/6}}} \right) \sim \frac{1}{k^{1/3}} \mathcal{F}(w) \quad , \quad \mathcal{F}(w) = \frac{1}{\pi} \int_0^\infty \text{Ai}^2(v - w) dv$$

Conclusion

- Connection between fermions in a rotating trap and the complex Ginibre ensemble of RMT
- « Wedding/birthday cake » structure as the rotating frequency decreases and higher Landau levels get populated
- Interesting edge behaviors and connection to the Airy kernel for very high Landau levels

Conclusion

- Connection between fermions in a rotating trap and the complex Ginibre ensemble of RMT
- « Wedding/birthday cake » structure as the rotating frequency decreases and higher Landau levels get populated
- Interesting edge behaviors and connection to the Airy kernel for very high Landau levels
- Study of the full counting statistics Smith, Le Doussal, Majumdar, G. S. '22
see also Charlier et al. '20-'22, Akemann, Byun, Ebke '22

Conclusion

- Connection between fermions in a rotating trap and the complex Ginibre ensemble of RMT
- « Wedding/birthday cake » structure as the rotating frequency decreases and higher Landau levels get populated
- Interesting edge behaviors and connection to the Airy kernel for very high Landau levels
- Study of the full counting statistics Smith, Le Doussal, Majumdar, G. S. '22
see also Charlier et al. '20-'22, Akemann, Byun, Ebke '22
- Extension to finite temperature
Kulkarni, Le Doussal, Majumdar, G. S. to appear soon

Conclusion

- Connection between fermions in a rotating trap and the complex Ginibre ensemble of RMT
- « Wedding/birthday cake » structure as the rotating frequency decreases and higher Landau levels get populated
- Interesting edge behaviors and connection to the Airy kernel for very high Landau levels
- Study of the full counting statistics Smith, Le Doussal, Majumdar, G. S. '22
see also Charlier et al. '20-'22, Akemann, Byun, Ebke '22
- Extension to finite temperature
Kulkarni, Le Doussal, Majumdar, G. S. to appear soon
- Generalization to other types of potentials



$$\frac{1}{2}\omega^2 r^2 \longrightarrow \frac{1}{2}\omega^2 r^2 + v\left(\frac{r}{\sqrt{N}}\right) \quad , \quad \text{e.g.} \quad v(x) = \frac{\gamma}{x^2}$$

■ Generalization to other types of potentials


Kulkarni, Majumdar, G. S. '21

Smith, Le Doussal, Majumdar, G. S. '22

$$\frac{1}{2}\omega^2 r^2 \longrightarrow \frac{1}{2}\omega^2 r^2 + v\left(\frac{r}{\sqrt{N}}\right) \quad , \quad \text{e.g.} \quad v(x) = \frac{\gamma}{x^2}$$




Happy Birthday Yan !



Quantum Fermi gas microscope

- Direct imaging of spatial fluctuations of the positions of fermions



M. Greiner et al., PRL 2015