


Tropical geometry for Nagata's conjecture and Legendrian curves

Nikita Kalinin

University of Geneva

December 8, 2015

Advocating

Tropical geometry has been recently heavily used in

- the proof of longstanding Rota log-concavity conjecture [Adiprasito, Huh, Katz, 2015]. In particular it implies that the coefficients a_i of the chromatic polynomial of a graph form a log-concave sequence, i.e. $a_i^2 \geq a_{i-1}a_{i+1}$.
- a uniform estimate for the number $N(d, g)$ of rational points on curves of small Mordell-Weil rank [Katz, Rabinoff, Zureick-Brown, 2015], as well as unconditional estimate for the number of torsion points of the images of curves in Jacobians.



Sandpile models: emergence of tropical objects

Topplings


Definition

A sandpile is a collection of indistinguishable sand grains distributed among a finite subset Γ of $\mathbb{Z}^2$, that is a function $f : \Gamma \to \mathbb{N}_0$. A vertex v is unstable if $f(v) \geq 4$. An unstable vertex can topple by sending one grain of sand to each of 4 neighbours.

Topplings

Definition

A sandpile is a collection of indistinguishable sand grains distributed among a finite subset Γ of $\mathbb{Z}^2$, that is a function $f : \Gamma \to \mathbb{N}_0$. A vertex v is unstable if $f(v) \geq 4$. An unstable vertex can topple by sending one grain of sand to each of 4 neighbours.



The problem: picture after relaxation

Definition

The process of doing topplings while it is possible is called relaxation.

Proposition

The order of topplings has no influence on the result of the relaxation.

The problem: picture after relaxation

Definition

The process of doing topplings while it is possible is called relaxation.

Proposition


The order of topplings has no influence on the result of the relaxation.

Definition


We chose a boundary $\partial\Gamma \subset \Gamma$, where we never do topplings. For example, $\Gamma = [0, \dots, N] \times [0, \dots, N]$, $\partial\Gamma$ is the border of this square.

We start from the maximal stable distribution of sand, i.e. $f \equiv 3$, then we add a grain somewhere. What is the result of relaxation? And if we add a number of grains to different places?


Simulation, results



Just after beginning...



... in the middle...



Et voilà ! Final result!

White: 3 grains, Green: 2 grains, Yellow: 1 grain, Red: no grains,
Black: more than 3 grains.

Explanation: look at the number of topplings

Consider the number $h(x, y)$ of topplings at a point (x, y) during the relaxation.

Proposition

If the number of sand grains at (x, y) after relaxation is the same as before the relaxation, then
$h(x-1, y) + h(x+1, y) + h(x, y-1) + h(x, y+1) - 4h(x, y) = 0.$

Explanation: look at the number of topplings

Consider the number $h(x, y)$ of topplings at a point (x, y) during the relaxation.


Proposition

If the number of sand grains at (x, y) after relaxation is the same as before the relaxation, then
$h(x-1, y) + h(x+1, y) + h(x, y-1) + h(x, y+1) - 4h(x, y) = 0$.

Proof.

Indeed, count the number of incoming and outgoing grains. □

That is, h is harmonic at (x, y). In fact, $h(x, y)$ is a "piece-wise" linear function.



Tropical geometry: explanation of pictures

Tropical curves

Definition

Let $\mathcal{A}$ be a finite subset of $\mathbb{Z}^2$. A tropical polynomial is a function $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}} (a_{ij} + ix + jy), a_{ij} \in \mathbb{R}$. Note that $h : \mathbb{R}^2 \to \mathbb{R}$.

Tropical curves

Definition

Let $\mathcal{A}$ be a finite subset of $\mathbb{Z}^2$. A tropical polynomial is a function $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}} (a_{ij} + ix + jy), a_{ij} \in \mathbb{R}$. Note that $h : \mathbb{R}^2 \to \mathbb{R}$.

A tropical curve is the set of non-smooth points of a tropical polynomial.

Tropical curves

Definition

Let $\mathcal{A}$ be a finite subset of $\mathbb{Z}^2$. A tropical polynomial is a function $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}} (a_{ij} + ix + jy), a_{ij} \in \mathbb{R}$. Note that $h : \mathbb{R}^2 \to \mathbb{R}$.

A tropical curve is the set of non-smooth points of a tropical polynomial.

Motivation: let $log_t(x,y) = (\log_t(|x|), \log_t(|y|))$, $log_t : \mathbb{C}^2 \to \mathbb{R}^2$.

Tropical curves

Definition

Let $\mathcal{A}$ be a finite subset of $\mathbb{Z}^2$. A tropical polynomial is a function $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}}(a_{ij} + ix + jy), a_{ij} \in \mathbb{R}$. Note that $h : \mathbb{R}^2 \to \mathbb{R}$.

A tropical curve is the set of non-smooth points of a tropical polynomial.

Motivation: let $log_t(x,y) = (\log_t(|x|), \log_t(|y|))$, $log_t : \mathbb{C}^2 \to \mathbb{R}^2$.

Consider a family of polynomials $F_t(x,y) = \sum_{(i,j)\in\mathcal{A}}(t^{a_{ij}} + \dots)x^i y^j$.

Define the corresponding curves $C_t = \{(x,y)|F_t(x,y) = 0\} \subset \mathbb{C}^2$. An amoeba of C is $log_t C$.

Tropical curves

Definition


Let $\mathcal{A}$ be a finite subset of $\mathbb{Z}^2$. A tropical polynomial is a function $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}}(a_{ij} + ix + jy), a_{ij} \in \mathbb{R}$. Note that $h : \mathbb{R}^2 \to \mathbb{R}$.

A tropical curve is the set of non-smooth points of a tropical polynomial.

Motivation: let $log_t(x,y) = (\log_t(|x|), \log_t(|y|))$, $log_t : \mathbb{C}^2 \to \mathbb{R}^2$. Consider a family of polynomials $F_t(x,y) = \sum_{(i,j)\in\mathcal{A}}(t^{a_{ij}} + \dots)x^i y^j$. Define the corresponding curves $C_t = \{(x,y)|F_t(x,y) = 0\} \subset \mathbb{C}^2$. An amoeba of C is $log_t C$.

Proposition

Then $C = \lim_{t\to 0} log_t C_t$ *is the tropical curve defined by* $h(x,y) = \min\limits_{(i,j)\in\mathcal{A}}(a_{ij} + ix + jy)$.



Nagata's conjecture: a curve through singular points

Singular points

We consider a polynomial $F(x,y) = \sum a_{ij}x^i y^j$, $a_{ij} \in \mathbb{C}, 0 \leq i,j, i+j \leq d$ of degree d, it defines a curve C as its zero-set, $C = \{(x,y) | F(x,y) = 0\}$.

Definition

A point $p \in \mathbb{C}^2$ is of multiplicity m for C if all the partial derivatives of F up to order $m-1$ vanish at p.

Singular points

We consider a polynomial $F(x,y) = \sum a_{ij}x^i y^j$, $a_{ij} \in \mathbb{C}, 0 \leq i,j, i+j \leq d$ of degree d, it defines a curve C as its zero-set, $C = \{(x,y) | F(x,y) = 0\}$.

Definition

A point $p \in \mathbb{C}^2$ is of multiplicity m for C if all the partial derivatives of F up to order $m-1$ vanish at p.

Example

The point $(0,0)$ is of multiplicity

- 1 if C passes through $(0,0)$
- 2 if $(0,0)$ is a singular point of C
- m if $a_{ij} = 0$ for $i + j \leq m - 1$.

Dimension counting

As we see, the fact that a point p is of multiplicity m for a curve C imposes $\frac{m(m+1)}{2}$ linear conditions on the coefficients a_{ij}.

Dimension counting

As we see, the fact that a point p is of multiplicity m for a curve C imposes $\frac{m(m+1)}{2}$ linear conditions on the coefficients a_{ij}.

A curve of degree d has $\frac{(d+1)(d+2)}{2}$ coefficients. Therefore, if $p_1, p_2, \ldots, p_n$ are in general position, a curve C passes through $p_1, p_2, \ldots, p_n$ with some multiplicities $m_1, m_2, \ldots m_n$, then we should expect

$$\frac{(d+1)(d+2)}{2} - 1 \geq \sum_{1 \leq i \leq n} \frac{m_i(m_i+1)}{2}.$$

Dimension counting

As we see, the fact that a point p is of multiplicity m for a curve C imposes $\frac{m(m+1)}{2}$ linear conditions on the coefficients a_{ij}.
A curve of degree d has $\frac{(d+1)(d+2)}{2}$ coefficients. Therefore, if $p_1, p_2, \ldots, p_n$ are in general position, a curve C passes through $p_1, p_2, \ldots, p_n$ with some multiplicities $m_1, m_2, \ldots m_n$, then we should expect

$$\frac{(d+1)(d+2)}{2} - 1 \geq \sum_{1 \leq i \leq n} \frac{m_i(m_i+1)}{2}.$$

Unfortunately, these linear conditions are **not** independent.

- A line (d=1) passes though 2 points. $\frac{2 \cdot 3}{2} - 1 \geq \frac{1 \cdot 2}{2} + \frac{1 \cdot 2}{2}$ (still true).
- A doubled conic ($d = 4$) passes through 5 points with multiplicities 2. $\frac{5 \cdot 6}{2} - 1 \geq 5 \cdot \frac{2 \cdot 3}{2}$ (not true).

Nagata's conjecture

As a reasonable estimate, Nagata conjectured that if $n \geq 10$ and the above conditions are satisfied, then $d > \frac{\sum m_i}{\sqrt{n}}$.

Nagata's conjecture

As a reasonable estimate, Nagata conjectured that if $n \geq 10$ and the above conditions are satisfied, then $d > \frac{\sum m_i}{\sqrt{n}}$.

- Open even for $n = 10, m_i = m$. (That is $d > \sqrt{n} \cdot m$)

Nagata's conjecture

As a reasonable estimate, Nagata conjectured that if $n \geq 10$ and the above conditions are satisfied, then $d > \frac{\sum m_i}{\sqrt{n}}$.

- Open even for $n = 10, m_i = m$. (That is $d > \sqrt{n} \cdot m$)
- Nagata proved it himself for $n = k^2$. The proof gives a counterexample for 14th Hilbert Problem:
  Is the ring of invariants of an algebraic group acting on a polynomial ring is always finitely generated?

Duality between tropical curves and subdivisions of Δ

A tropical curve comes as the locus where a function $h(x, y) = \max\limits_{(i,j)\in\Delta} (a_{ij} + ix + jy)$ is not smooth.

Duality between tropical curves and subdivisions of Δ

A tropical curve comes as the locus where a function $h(x,y) = \max\limits_{(i,j)\in\Delta}(a_{ij} + ix + jy)$ is not smooth.

Definition

The Extended Newton polytope of h is the convex hull of the set $\bigcup\{(i,j,s)|(i,j) \in \Delta, s \leq a_{ij}\}$.

Duality between tropical curves and subdivisions of Δ

A tropical curve comes as the locus where a function $h(x,y) = \max\limits_{(i,j)\in\Delta}(a_{ij} + ix + jy)$ is not smooth.

Definition

The Extended Newton polytope of h is the convex hull of the set $\bigcup\{(i,j,s)|(i,j) \in \Delta, s \leq a_{ij}\}$.

The projection along the third coordinate of the faces of the Extended Newton polygon gives a subdivision of the convex hull of Δ.

Duality between tropical curves and subdivisions of Δ

A tropical curve comes as the locus where a function $h(x,y) = \max\limits_{(i,j)\in\Delta}(a_{ij} + ix + jy)$ is not smooth.

Definition


The Extended Newton polytope of h is the convex hull of the set $\bigcup\{(i,j,s)|(i,j) \in \Delta, s \leq a_{ij}\}$.

The projection along the third coordinate of the faces of the Extended Newton polygon gives a subdivision of the convex hull of Δ.


Proposition

This subdivision is dual to the tropical curve C defined by h.


Example



(A)



(B)



(C)

The extended Newton polyhedron of the curve C is drawn in (A). The projection of its faces gives us the subdivision of the Newton polygon of the curve, see (B). The tropical curve C is drawn in (C).

Influenced regions

Definition

For a point P on a tropical curve C we define the region of influence of P in the Newton polygon as follows. Draw **straight lines** through all the edges of C which pass through P. Take **all the vertices** of C lying on the latter lines. Take **the dual cells** to these vertices. The union of these cells is the region of influence of P.


Influenced regions


Definition


For a point P on a tropical curve C we define the region of influence of P in the Newton polygon as follows. Draw **straight lines** through all the edges of C which pass through P. Take **all the vertices** of C lying on the latter lines. Take **the dual cells** to these vertices. The union of these cells is the region of influence of P.


On the previous slide P is the tropicalization of a singular point of multiplicity 3. The influenced region is the whole Newton polygon.

Examples









Theorems

Theorem

If P is of multiplicity m on C, then the area of the influenced region of the Newton polygon is bigger or equal than

- $\frac{3}{8}m^2$ *if* $\mathrm{Val}(P)$ *is a vertex of* $\mathrm{Trop}(C)$
- $\frac{1}{2}m^2$ *if* $\mathrm{Val}(P)$ *is not a vertex of* $\mathrm{Trop}(C)$.

Theorems

Theorem

If P is of multiplicity m on C, then the area of the influenced region of the Newton polygon is bigger or equal than

- $\frac{3}{8}m^2$ *if* $\mathrm{Val}(P)$ *is a vertex of* $\mathrm{Trop}(C)$
- $\frac{1}{2}m^2$ *if* $\mathrm{Val}(P)$ *is not a vertex of* $\mathrm{Trop}(C)$.

Theorem

If points $P_1, P_2, \ldots P_n$ are chosen generically, then all triple intersections of their regions of influence are empty.


Implications

These tropical theorems imply some estimates in Nagata's conjecture.

Proposition

The degree d of a curve passing through generic $p_1, p_2, \ldots, p_n$ with multiplicities m is subject to the following inequality:

$$d^2 \geq (n - \sqrt{n} - 1)m^2$$



Tropical differential forms in Legendrian geometry.

Holomorphic Legendrian curves

Definition

A differential 1-form ω on M^3 is contact if $\omega \wedge d\omega \neq 0$ everywhere.

Holomorphic Legendrian curves

Definition

A differential 1-form ω on M^3 is contact if $\omega \wedge d\omega \neq 0$ everywhere.

We can take the same definition for holomorphic forms, the only complex contact form on $\mathbb{C}P^3$ is $\omega = ydx - xdy + wdz - zdw$.

Holomorphic Legendrian curves

Definition

A differential 1-form ω on M^3 is contact if $\omega \wedge d\omega \neq 0$ everywhere.

We can take the same definition for holomorphic forms, the only complex contact form on $\mathbb{C}P^3$ is $\omega = ydx - xdy + wdz - zdw$.

Definition

A curve C is contact or Legendrian if $\omega(TC) = 0$.

Holomorphic Legendrian curves

Definition

A differential 1-form ω on M^3 is contact if $\omega \wedge d\omega \neq 0$ everywhere.

We can take the same definition for holomorphic forms, the only complex contact form on $\mathbb{C}P^3$ is $\omega = ydx - xdy + wdz - zdw$.

Definition

A curve C is contact or Legendrian if $\omega(TC) = 0$.

How many algebraic Legendrian curves of a given degree pass through given points in $\mathbb{C}P^3$?

Holomorphic Legendrian curves

Definition

A differential 1-form ω on M^3 is contact if $\omega \wedge d\omega \neq 0$ everywhere.

We can take the same definition for holomorphic forms, the only complex contact form on $\mathbb{C}P^3$ is $\omega = ydx - xdy + wdz - zdw$.

Definition

A curve C is contact or Legendrian if $\omega(TC) = 0$.

How many algebraic Legendrian curves of a given degree pass through given points in $\mathbb{C}P^3$?
Answer: **three** rational Legendrian cubics through 3 points and one line.

Exact formulae

All the curves tangent to $ydx - xdy + wdz - zdw$ and passing through three points $(0,0,0,1), (1,1,1,1), (-1,1-1,1) \in \mathbb{C}P^3$ are:
$(3t - t^3, 2t^2 + 2\mu(t - t^3), 2t^3, 1 + t^2 - 2\mu(t - t^3)), t \in \mathbb{C}P^1, \mu \in \mathbb{C}$.

Exact formulae

All the curves tangent to $ydx - xdy + wdz - zdw$ and passing through three points $(0,0,0,1), (1,1,1,1), (-1,1-1,1) \in \mathbb{C}P^3$ are:
$(3t - t^3, 2t^2 + 2\mu(t - t^3), 2t^3, 1 + t^2 - 2\mu(t - t^3)), t \in \mathbb{C}P^1, \mu \in \mathbb{C}$.

Question

What is the compactification of this family? ($\mu \to \infty$)

Exact formulae

All the curves tangent to $ydx - xdy + wdz - zdw$ and passing through three points $(0,0,0,1), (1,1,1,1), (-1,1-1,1) \in \mathbb{C}P^3$ are:
$(3t - t^3, 2t^2 + 2\mu(t - t^3), 2t^3, 1 + t^2 - 2\mu(t - t^3)), t \in \mathbb{C}P^1, \mu \in \mathbb{C}$.

Question

What is the compactification of this family? ($\mu \to \infty$)

Answer: three lines in a plane, with intersection at one point. This corresponds to the degenerate constraints: three points and a line in one plane.

Exact formulae

All the curves tangent to $ydx - xdy + wdz - zdw$ and passing through three points $(0,0,0,1), (1,1,1,1), (-1,1-1,1) \in \mathbb{C}P^3$ are:
$(3t - t^3, 2t^2 + 2\mu(t - t^3), 2t^3, 1 + t^2 - 2\mu(t - t^3)), t \in \mathbb{C}P^1, \mu \in \mathbb{C}$.

Question


What is the compactification of this family? ($\mu \to \infty$)

Answer: three lines in a plane, with intersection at one point. This corresponds to the degenerate constraints: three points and a line in one plane.

Question

Is it true that there are at least d rational legendrian curves of degree d through generic d points in $\mathbb{C}P^3$ and a line?

Tropical Legendrian curves



The tropicalization of the surface swept by legendrian cubics through three points. A particular tropical curve inside.

Abstract tropical curves

Proposition

Note that the amoeba $\log_t(C)$ *is given by real part of the integrals of* $\frac{dx}{x}, \frac{dy}{y}$ *on* C *and rescaling by* $\ln t$.

Abstract tropical curves

Proposition

Note that the amoeba $\log_t(C)$ *is given by real part of the integrals of* $\frac{dx}{x}, \frac{dy}{y}$ *on* C *and rescaling by* $\ln t$.

Definition (see more details in the thesis of Lionel Lang)

A tropical curve S is a degeneration of a family $S_t, t \to \infty$ of complex surfaces if the asymptotics of the lengths of the geodesics in the pair-of-pants decomposition are $\frac{1}{a_i \ln t}$. The corresponding edges of S have lengths a_i.

The degeneration comes with the contraction maps $S_t \to S$, where we contract the hyperbolic collars of the geodesics to the edges of a tropical curve.

Tropical differential forms

Definition

We define tropical differential form by the following family.

Tropical differential forms

Definition

We define tropical differential form by the following family. We have a family S_t of **punctured** Riemann surfaces, that degenerates to a tropical curve S.

Tropical differential forms

Definition

We define tropical differential form by the following family. We have a family S_t of **punctured** Riemann surfaces, that degenerates to a tropical curve S. Also we have a family ω_t of differentials with poles of 3rd kind (i.e. the residue is not zero) at the punctures.

Tropical differential forms

Definition


We define tropical differential form by the following family. We have a family S_t of **punctured** Riemann surfaces, that degenerates to a tropical curve S. Also we have a family ω_t of differentials with poles of 3rd kind (i.e. the residue is not zero) at the punctures.

Proposition

If $\lim \int_{\gamma_t} \omega_t = L$ *then* $\lim \frac{1}{\ln t} \int_{\Gamma_t} \omega_t = aL$. *So,* $\lim \omega$ *should be though as* $L \cdot dX$

(Here γ_t is the geodesic in the pair-of-pants decomposition, Γ_t is a path in the collar from one boundary component to another, a is the length of the corresponding tropical edge.)

Application of tropical differential forms



A part of the tropicalization of a legendrian curve. This part looks like a tropical line.

We can prove that the plane $x + y = z$ divides the interval between two vertices in the middle.

Summary

- Sandpile models and their evolution.

Summary

- Sandpile models and their evolution.
- For the square in $\mathbb{Z}^2$ the vertices with less than 3 grains constitute a tropical curve.

Summary

- Sandpile models and their evolution.
- For the square in $\mathbb{Z}^2$ the vertices with less than 3 grains constitute a tropical curve.
- From the matroidal point of view, tropical geometry is about subdivisions of the Newton polygon of a curve.

Summary

- Sandpile models and their evolution.
- For the square in $\mathbb{Z}^2$ the vertices with less than 3 grains constitute a tropical curve.
- From the matroidal point of view, tropical geometry is about subdivisions of the Newton polygon of a curve.
- Constraints, imposed by singular points on the coefficients of a curve, are visible on this subdivision.

Summary

- Sandpile models and their evolution.
- For the square in $\mathbb{Z}^2$ the vertices with less than 3 grains constitute a tropical curve.
- From the matroidal point of view, tropical geometry is about subdivisions of the Newton polygon of a curve.
- Constraints, imposed by singular points on the coefficients of a curve, are visible on this subdivision.
- A definition of an abstract tropical curve and a tropical differential form.


Thank you for your attention!
Geneva, 2015