

# PEDIATRICS

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## COVID-19 in Children and the Dynamics of Infection in Families

Klara M. Posfay-Barbe, MD, Noemie Wagner, MD, Magali Gauthey, MD,  
Dehlia Moussaoui, MD, Natasha Loevy, MD, Alessandro Diana, MD, Arnaud G. L'Huillier, MD

**DOI:** 10.1542/peds.2020-1576

**Journal:** *Pediatrics*

**Article Type:** Research Brief

**Citation:** Posfay-Barbe KM, Wagner N, Gauthey M, et al. COVID-19 in children and the dynamics of infection in families. *Pediatrics*. 2020; doi: 10.1542/peds.2020-1576

This is a pre-publication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

## COVID-19 in Children and the Dynamics of Infection in Families

Klara M. Posfay-Barbe, MD<sup>1</sup>, Noemie Wagner, MD<sup>1</sup>, Magali Gauthey, MD<sup>2</sup>, Dehlia Moussaoui, MD<sup>3</sup>, Natasha Loevy, MD<sup>4</sup>, Alessandro Diana, MD<sup>5,6</sup>, Arnaud G. L'Huillier, MD<sup>1,7</sup>

<sup>1</sup>Pediatric Infectious Diseases Unit, Department of Woman, Child and Adolescent Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Geneva, Switzerland

<sup>2</sup>Pediatric Department, Hôpital de La Tour, Geneva, Switzerland

<sup>3</sup>Division of General Pediatrics, Department of Woman, Child and Adolescent Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Geneva, Switzerland

<sup>4</sup>Pediatric Platform for Clinical Research, Department of Woman, Child and Adolescent Medicine, Geneva University Hospitals and Faculty of Medicine, University of Geneva, Geneva, Switzerland

<sup>5</sup>Lecturer, Primary Care Unit, University of Geneva, Geneva, Switzerland

<sup>6</sup>Clinique des Grangettes, Geneva, Switzerland

<sup>7</sup>Division of Infectious Diseases and Laboratory of Virology, Division of Laboratory Medicine, Geneva University Hospitals and Medical School, Geneva, Switzerland

### Corresponding author:

Arnaud G. L'Huillier, MD

Department of Woman, Child and Adolescent Medicine, Geneva University Hospitals and Medical School, Geneva, Switzerland Pediatric Infectious Diseases Unit

6 rue Willy-Donze, 1211 Geneva 14, Switzerland

+41 79 55 31 385; [arnaud.lhuillier@hcuge.ch](mailto:arnaud.lhuillier@hcuge.ch)

**Financial disclosure:** The authors have no financial relationships relevant to this article to disclose.

**Funding source:** no funding

**Potential Conflicts of Interest:** The authors have no conflicts of interest relevant to this article to disclose.

**Abbreviations:**

COVID-19: coronavirus disease; HHC: household contact; IQR: interquartile range; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2

**Contributors' Statement:** Klara Posfay-Barbe, Noemie Wagner and Arnaud G. L'Huillier conceived and designed the study, designed the data collection instruments, coordinated and supervised data collection, carried out the initial analyses, drafted the initial manuscript, and reviewed and revised the manuscript. Magali Gauthey, Dehlia Moussaoui, Natasha Loevy and Alessandro Diana coordinated and supervised data collection, critically reviewed the manuscript for important intellectual content, and reviewed and revised the manuscript. All the authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

## **Introduction**

Since the onset of coronavirus disease (COVID-19) pandemic, children have been less affected than adults in terms of severity<sup>1-3</sup> and frequency, accounting for <2% of the cases<sup>2-5</sup>. Unlike with other viral respiratory infections, children do not seem to be a major vector of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transmission, with most pediatric cases described inside familial clusters<sup>6</sup> and no documentation of child-to-child or child-to-adult transmission<sup>7,8</sup>. The aim of this work was to describe the clinical presentation of the first 40 pediatric cases of COVID-19 in our city and the dynamics of their familial clusters.

## **Methods**

From March 10<sup>th</sup> to April 10<sup>th</sup> 2020, all patients <16 years old with SARS-CoV-2 infection were identified by means of the Geneva University Hospital's surveillance network (Switzerland). The network notifies the institution's pediatric infectious diseases specialists about results of nasopharyngeal specimens tested for SARS-CoV-2 by reverse-transcription polymerase-chain-reaction. This study was approved by the Regional Ethics Committee. After informed oral parental consent and its documentation in the medical charts, chart reviews were used to retrieve clinical data and parents were called for patients and household contacts (HHC) follow-up. HHC were considered suspect if they had fever or acute respiratory symptoms, as per the Swiss Federal Office for Public Health's case definition<sup>9</sup> (Supplementary Methods).

Categorical data were compared using the chi-squared test, with p-values <0.05 considered significant. Statistics were performed using SPSS version 23.0.

## **Results**

Among a total of 4310 SARS-CoV-2 cases, 40 were <16 years old (0.9%). One patient for which telephone follow-up was not possible was excluded because of the inability to evaluate clinical evolution and household contact (HHC) symptoms. The median follow-up of the households was 18 days (interquartile range [IQR] 14-28).

**Clinical presentation, diagnosis and management.** Demographics, clinical presentation and diagnosis of the study children are detailed in Table 1. Of note, 29 (74%) patients were previously healthy; the most frequently reported comorbidities were asthma (10%), diabetes (8%), obesity (5%), premature birth (5%) and hypertension (3%). Seven patients (18%) were hospitalized to the ward, for a median duration of 3 days (IQR 2-4); reasons for admission were surveillance for non-hypoxemic viral pneumonia (n=2), fever without source (n=2), apparent life-threatening event (n=1), sepsis-like event (n=1) and one paucisymptomatic child admitted because both parents had severe COVID-19 (n=1). No patient required intensive care unit admission or SARS-CoV-2 specific therapies. The others 32 patients were managed as outpatients. All patients had a complete resolution of symptoms by day seven after diagnosis.

**Familial clusters.** Familial cluster evaluation revealed a t number of four household members per family (IQR 3-4). Among the 111 HHC of study children, mothers predominated (n=39), followed by fathers (n=32), pediatric siblings (n=23), adult siblings (n=8) and grand-parents (n=7) (Figure 1). Adult HHC were suspected or confirmed with COVID-19 prior to the study child in 79% (31/39) of cases. In only 8% (3/39) of households did the study child develop symptoms prior to any other HHC (Figure 1). Interestingly, 85% (75/88) of adult HHC developed symptoms at some point, compared to 43% (10/23) of pediatric HHC (p<0.001). Also, 92% (36/39) of mothers developed symptoms, compared to 75% (24/32) of fathers (p=0.04).

## **Discussion**

Most children in our study had mild or atypical presentations: headache and nasal discharge were described in more than half of cases, and anosmia and abdominal symptoms in more than 20%, which is more frequent than previously described<sup>2</sup>. Some of these symptoms might be underreported because younger children may not be able to describe them.

In 79% of households, at least one adult family member was suspected or confirmed for COVID-19 prior to symptom onset in the study child, confirming that children are infected mainly inside familial clusters<sup>6</sup>. Surprisingly, in 33% of households, symptomatic HHC tested negative despite belonging to a familial cluster with confirmed SARS-CoV-2 cases, suggesting an underreporting of cases. In only 8% of households did a child develop symptoms prior to any other HHC, which is in line with previous data showing that children are index cases in less than 10% of SARS-CoV-2 familial clusters<sup>10</sup>; however, our study design cannot confirm that child-to-adult transmission occurred.

This study has some limitations. The study sample likely doesn't represent the total number of pediatric SARS-CoV-2 cases during this time period. Indeed, patients with milder or atypical presentation might not have sought medical attention. Moreover, the recall of symptom onset among HHC might be inaccurate, even though this seems for once less likely due to the confinement measures and anxiety in the community.

The results of this study are important because of the extensive household contact tracing and the almost absence of loss to follow-up. Extended diagnostic screening of suspected cases and thorough contact tracing are needed to better understand the dynamics of transmission within households.

## References

1. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6):e20200702.
2. Center for Disease Control and Prevention (CDC) COVID response team. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):422-426.
3. Livingston E, Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy [published online March 17, 2020]. *JAMA*. 2020. doi: 10.1001/jama.2020.4344.
4. Tagarro A, Epalza C, Santos M, et al. Screening and Severity of Coronavirus Disease 2019 (COVID-19) in Children in Madrid, Spain [published online April 8, 2020]. *JAMA Pediatr*. doi:10.1001/jamapediatrics.2020.1346
5. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239-1242.
6. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children [published online March 18, 2020]. *N Engl J Med*. doi: 10.1056/NEJMc200507
7. World Health Organization. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). Published February 28, 2020.
8. Danis K, Epaulard O, Benet T, et al. Cluster of coronavirus disease 2019 (Covid-19) in the French Alps, 2020 [published online April 11, 2020]. *Clin Infect Dis*. doi: 10.1093/cid/ciaa424
9. Federal Office of Public Health. Accessed April 22, 2020 [Available from: <https://www.bag.admin.ch/bag/en/home.html>].
10. Zhu Y, Bloxham CJ, Hulme KD, et al. Children are unlikely to have been the primary source of household SARS-CoV-2 infections [published online March 30, 2020]. *medRxiv*. doi:2020.2003.2026.20044826.

Table 1. Demographics and clinical presentation of study patients

	<b>n=39</b>
<b><u>Demographics</u></b>	
Median age, years (IQR)	11.1 (5.7-14.5)
Female gender, No. (%)	22 (56)
<b><u>Clinical presentation</u></b>	
<u>Median time between symptom onset and diagnosis, days (IQR)</u>	2 (1-3)
<u>Reported symptoms, No. (%)</u>	
Cough	32 (82)
Fever	26 (67)
Nasal discharge	25 (64)
Headache	22 (56)
Sore throat	14 (36)
Shortness of breath	13 (33)
Myalgia	13 (33)
Abdominal pain	11 (28)
Anosmia	8 (21)
Arthralgia	7 (18)
Diarrhea	7 (18)
Fatigue	5 (13)
Rash	5 (13)
Dysgeusia	4 (10)
Nausea	4 (10)
Vomiting	3 (8)
Thoracic pain	2(5)
Conjunctivitis	1 (3)
<b><u>Diagnosis, No. (%)</u></b>	
Upper respiratory tract infection	27 (69)
Influenza-like illness	2(5)
Fever without source	2(5)

## Pre-publication Release

Pneumonia	2(5)
Obstructive bronchitis	2(5)
Sepsis-like event	1 (3)
Croup	1 (3)
ALTE	1 (3)
Asymptomatic	1 (3)

---

IQR: interquartile range; ALTE: Apparent life-threatening events

Figure 1. Description of individual household clusters with asymptomatic, suspected and confirmed SARS-CoV-2 cases

Patient →	1	2	3	4	5	6	7	8	9	10	11	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	Total	Symptomatic			
RT-PCR CT value	16	22	21	22	30	34	34	19	32	33	26	31	16	16	14	17	24	33	16	19	13	28	20	23	24	38	24	32	34	37	34	28	35	32	31	35	36	32	36	n	n (%)			
Mother	+			+	+	-				+	-					-	+	+	+	+	+	+	+	+		+			-	+	+		-	+	+	+	-	-		39	36 (92)			
Father	-			+		+		+			+					+			+	+	+	+	+	+	+												+	+		32	24 (75)			
Adult Sibling 1																	+						+	+																				
Adult Sibling 2																																										8	8 (100)	
Grand-Parent 1							+									+																						+	+					
Grand-Parent 2																+																										7	5 (71)	
Other Adult																																										2	2 (100)	
Pediatric Sibling 1					-					-																																		
Pediatric Sibling 2																																												
Pediatric Sibling 3																																												
Pediatric Sibling 4																																											23	10 (43)

Green, yellow and red squares correspond to symptomatic household contacts who developed symptoms respectively, before, simultaneously and after study patients. White squares correspond to asymptomatic household contacts. The “+” and “-” signs correspond to the results of SARS-CoV-2 nasopharyngeal RT-PCR; patients without testing have an empty square. The absence of a family member inside a given household cluster is shown in grey.

The study patient was the first to develop symptoms in clusters #2, #3 and #13 only.

CT values correspond to the number of PCR cycles required to amplify the virus; hence the CT value is inversely proportional to the viral load.

SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2; RT-PCR: reverse-transcription polymerase chain reaction; CT: cycle threshold.

### **Supplementary Methods**

Study setting. The study was performed while schools, daycares, restaurants, bars and shops were closed. Citizens were allowed to circulate outside their housing in groups of five or less while respecting social distance measures.

RT-PCR testing criteria. For epidemiological purposes and to maximize contact tracing, every patient with respiratory symptoms or fever who came to the hospital was tested for SARS-CoV-2 by RT-PCR, independently of epidemiological links. There was no significant shortage of testing capacity in our institution during this time period. The data and results of household contacts (HHC) RT-PCR testing was available to the study team through the hospital's surveillance network.

Definitions. HHCs were defined as persons living in the same housing as the SARS-CoV-2 RT-PCR positive study patient. HHC were asked whether they developed symptoms before, after or at the same time as the study patient; no minimal threshold was used for this definition.

## COVID-19 in Children and the Dynamics of Infection in Families

Klara M. Posfay-Barbe, Noemie Wagner, Magali Gauthey, Dehlia Moussaoui, Natasha Loevy, Alessandro Diana and Arnaud G. L'Huillier  
*Pediatrics* originally published online May 26, 2020;

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/early/2020/05/22/peds.2020-1576.citation>

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **COVID-19 in Children and the Dynamics of Infection in Families**

Klara M. Posfay-Barbe, Noemie Wagner, Magali Gauthey, Dehlia Moussaoui, Natasha Loevy, Alessandro Diana and Arnaud G. L'Huillier  
*Pediatrics* originally published online May 26, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/05/22/peds.2020-1576.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

