

CYLINDRIC PARTITIONS AND SOME NEW A_2 ROGERS–RAMANUJAN IDENTITIES

SYLVIE CORTEEL, JEHANNE DOUSSE, AND ALI KEMAL UNCU

ABSTRACT. We study the generating functions for cylindric partitions with profile (c_1, c_2, c_3) for all c_1, c_2, c_3 such that $c_1 + c_2 + c_3 = 5$. This allows us to discover and prove seven new A_2 Rogers–Ramanujan identities modulo 8 with quadruple sums, related with work of Andrews, Schilling, and Warnaar.

1. INTRODUCTION AND STATEMENT OF RESULTS

A partition of a positive integer n is a non-increasing sequence of natural numbers whose sum is n . For example, the partitions of 4 are $4, 3 + 1, 2 + 2, 2 + 1 + 1$, and $1 + 1 + 1 + 1$.

The Rogers–Ramanujan identities [RR19], given by

$$(1.1) \quad \sum_{n \geq 0} \frac{q^{n^2}}{(q; q)_n} = \frac{1}{(q; q^5)_\infty (q^4; q^5)_\infty},$$

$$(1.2) \quad \sum_{n \geq 0} \frac{q^{n^2+n}}{(q; q)_n} = \frac{1}{(q^2; q^5)_\infty (q^3; q^5)_\infty},$$

are probably the most famous identities in the theory of q -series and partitions. Here we used the standard q -series notation, for $n \in \mathbb{N} \cup \{\infty\}$ and $j \in \mathbb{N}$,

$$(a; q)_n := \prod_{k=0}^{n-1} (1 - aq^k),$$

$$(a_1, \dots, a_j; q)_n := (a_1; q)_n \cdots (a_j; q)_n.$$

We also use the convention that $1/(q; q)_n = 0$ for negative n .

The Rogers–Ramanujan identities were interpreted in terms of partitions by MacMahon [Mac16] and Schur [Sch17] independently, and can be formulated as follows.

Theorem 1.1 (Rogers–Ramanujan identities, partition version). *Let $i = 0$ or 1 . For every natural number n , the number of partitions of n such that the difference between two consecutive parts is at least 2 and the part 1 appears at most i times is equal to the number of partitions of n into parts congruent to $\pm(2 - i) \pmod{5}$.*

In addition to Combinatorics and Number Theory, the Rogers–Ramanujan are related to several other fields of mathematics such as representation theory of affine Lie algebras [LW84, LW85], mathematical physics [Bax81], and algebraic geometry [BMS13], to name only a few. They were proved and generalized in many ways over the years, see for example [And74, And89, Bre79, Bre83, Cor17, GM81, Gor61, GOW16, Pas20]. The book of A. Sills give a good introduction to the subject [Sil17].

Among these generalizations, an important one is due to Gordon [Gor61], who embedded Theorem 1.1 in an infinite family of combinatorial identities.

Theorem 1.2 (Gordon’s identities). *Let r and i be integers such that $r \geq 2$ and $1 \leq i \leq r$. Let $S_{r,i}(n)$ be the number of partitions $\lambda = \lambda_1 + \lambda_2 + \cdots + \lambda_s$ of n such that $\lambda_j - \lambda_{j+r-1} \geq 2$ for all j , and at most $i - 1$ of the λ_j are equal to 1. Let $P_{r,i}(n)$ be the number of partitions of n whose parts are not congruent to $0, \pm i \pmod{2r + 1}$.*

2010 *Mathematics Subject Classification*. 05A15, 05A17, 05A30, 05E10, 11P81, 11P84.

Key words and phrases. cylindric partitions, A_2 Rogers–Ramanujan identities, q -series.

Then for every positive integer n , we have

$$S_{r,i}(n) = P_{r,i}(n).$$

The Rogers–Ramanujan identities correspond to the cases $r = i = 2$ and $r = 2, i = 1$.

Andrews later extended Gordon’s identities in a q -series form [And74], which can be stated as follows.

Theorem 1.3 (Andrews–Gordon identities). *Let $r \geq 2$ and $1 \leq i \leq r$ be two integers. We have*

$$(1.3) \quad \sum_{n_1 \geq \dots \geq n_{r-1} \geq 0} \frac{q^{n_1^2 + \dots + n_{r-1}^2 + n_i + \dots + n_{r-1}}}{(q; q)_{n_1}} \left[\begin{matrix} n_1 \\ n_1 - n_2 \end{matrix} \right]_q \dots \left[\begin{matrix} n_{r-2} \\ n_{r-2} - n_{r-1} \end{matrix} \right]_q = \frac{(q^{2r+1}, q^i, q^{2r-i+1}; q^{2r+1})_\infty}{(q; q)_\infty},$$

where for two integers n and m ,

$$\left[\begin{matrix} m+n \\ m \end{matrix} \right]_q := \begin{cases} \frac{(q; q)_{m+n}}{(q; q)_m (q; q)_n} & \text{for } m, n \geq 0, \\ 0 & \text{otherwise,} \end{cases}$$

is the classical q -binomial coefficient.

Note that (1.1) (resp. (1.2)) is the particular case of (1.3) where $r = i = 2$ (resp. $r = 2$ and $i = 1$).

The Andrews–Gordon identities can be proved using the powerful machinery of Bailey pairs, first introduced by Bailey [Bai49], later developed by Andrews [And84a] and generalised to the Bailey lattice by Agrawal, Andrews, and Bressoud [AAB87]. All the identities provable by using classical Bailey lattice are related to certain characters of the Lie algebra A_1 [ASW99], and are therefore called A_1 Rogers–Ramanujan identities. In the beginning of the 1990s, Milne and Lilly [ML92, ML95] extended the Bailey lemma to A_{n-1} for general n . However, even though their work led to many q -series identities, they did not discover any A_{n-1} Rogers–Ramanujan identities.

In 1999, Andrews, Schilling and Warnaar [ASW99] were able to describe an A_2 Bailey lemma and Bailey lattice. This allowed them to prove four A_2 Rogers–Ramanujan identities, one of which is the following:

Theorem 1.4 (Andrews–Schilling–Warnaar).

$$\sum_{n_1, n_2 \geq 0} \frac{q^{n_1^2 + n_2^2 - n_1 n_2}}{(q; q)_{n_1}} \left[\begin{matrix} 2n_1 \\ n_2 \end{matrix} \right]_q = \frac{1}{(q, q, q^3, q^4, q^6, q^6; q^7)_\infty}.$$

Shortly after, Warnaar [War06] showed how to prove these identities using Hall–Littlewood polynomials.

In their paper, Andrews, Schilling and Warnaar found several very general families of sum-product identities (Theorems 5.1, 5.3, and 5.4 of [ASW99]). They are quite intricate to state, so we will not do it in full generality here. However, let us note the important fact that, after multiplication by $(q; q)_\infty$, the product side of these identities can be interpreted in terms of characters of the W_3 algebra. However, in general, after multiplication by $(q; q)_\infty$, the sum-side does not have obviously positive coefficients anymore, even though we know the coefficients are positive because of their interpretation as characters. So far, the four A_2 Rogers–Ramanujan identities of Andrews, Schilling, and Warnaar (Theorem 1.4 and three other identities of the same shape) and a fifth identity of the same family conjectured by Feigin, Foda, and Welsh [FFW08] and proved by Corteel and Welsh [CW19] were the only cases in which the character could be expressed as a sum with obviously positive coefficients.

For example, the particular case $k = 3, i = 1$ in Theorem 5.3 of [ASW99] can be stated as follows.

Theorem 1.5 (Andrews–Schilling–Warnaar).

$$(q, q)_\infty \sum_{a_1, b_1, a_2, b_2 \in \mathbb{Z}} \frac{q^{a_1^2 + b_1^2 + a_2^2 + b_2^2 - a_1 b_1 + a_2 b_2 + a_1 + a_2 + b_1 + b_2}}{(q; q)_{a_1 - a_2} (q; q)_{b_1 - b_2} (q; q)_{a_2} (q; q)_{b_2} (q; q)_{a_2 + b_2 + 1}} = \frac{1}{(q^2, q^3, q^3, q^4, q^4, q^5, q^5, q^6; q^8)_\infty}$$

The product on the right represents a character, but the $(q, q)_\infty$ prevents us from interpreting the left-hand side as a sum with positive coefficients.

In 2016, Foda and Welsh [FW16] had the brilliant idea to prove the Andrews–Gordon identities and the Bressoud identities using the combinatorics of cylindric partitions. We will define these objects in Section 2.

Soon after, Corteel [Cor17] gave a combinatorial proof of the Rogers–Ramanujan identities using cylindric partitions and the Robinson–Schensted–Knuth correspondence. Corteel and Welsh in [CW19] built on these ideas to give a short proof of Andrews, Schilling and Warnaar’s four A_2 Rogers–Ramanujan identities and the fifth one conjectured by Feigin, Foda, and Welsh [FFW08], using the combinatorics of cylindric partitions as well.

Our goal in this paper is to continue these lines of work by presenting a new family of A_2 Rogers–Ramanujan identities using cylindric partitions. These new identities also give an expression of Andrews, Schilling and Warnaar’s products related to characters as sums with obviously positive coefficients.

We prove simultaneously seven new identities. The four simpler ones are stated below, while the three more complicated ones can be found in Theorem 2.5.

Theorem 1.6. *We have*

$$\begin{aligned} \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q &= \frac{1}{(q^2, q^3, q^3, q^4, q^4, q^5, q^5, q^6; q^8)_\infty}, \\ \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q &= \frac{1}{(q, q^2, q^3, q^4, q^4, q^5, q^6, q^7; q^8)_\infty}, \\ \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_3 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q &= \frac{1}{(q, q, q^3, q^3, q^5, q^5, q^7, q^7; q^8)_\infty}, \\ \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q &= \frac{1}{(q, q, q^2, q^4, q^4, q^6, q^7, q^7; q^8)_\infty}. \end{aligned}$$

The first equation of our theorem gives an expression of the product in Theorem 1.5 as a sum with obviously positive coefficients, and we give a combinatorial interpretation in terms of cylindric partitions. More generally, our full main result, Theorem 2.5, gives a positive sum expression for all the products in the case $k = 3$ in Theorem 5.3 of Andrews, Schilling, and Warnaar’s paper [ASW99].

The two challenges in proving Theorem 2.5 were the following: first, we “guessed” the sum-side by guessing the generating functions for certain cylindric partitions with bounded entries. Second, we proved these identities simultaneously by using a result of Corteel and Welsh (see Theorem 2.3). As it is quite intricate to handle quadruple sums, we used a recent computer algebra implementation of Ablinger and Uncu [AU19] and we give an automated proof of the main result.

This paper is organized as follows. In Section 2 we introduce cylindric partitions and the methodology. To give a simple example on how to use cylindric partitions to prove partitions identities, we prove the classical Rogers–Ramanujan identities. In Section 3, we prove Theorem 2.5.

Acknowledgements. JD wants to thank the hospitality of the department of Mathematics at UC Berkeley where some of the results were proven. JD is partially funded by the ANR COMBINÉ ANR-19-CE48-0011. The research of AU is supported by the Austrian Science Fund (FWF) SFB50-11 Project. The authors thank Ole Warnaar and Trevor Welsh for their interest and helpful remarks.

2. INTRODUCTION AND METHODOLOGY

Cylindric partitions were introduced by Gessel and Krattenthaler [GK97]. They are defined as follows.

Definition. Let k and ℓ be positive integers. Let $c = (c_1, c_2, \dots, c_k)$ be a composition, where $c_1 + c_2 + \dots + c_k = \ell$. A *cylindric partition with profile c* is a vector partition $\Lambda = (\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$, where each $\lambda^{(i)} = \lambda_1^{(i)} + \lambda_2^{(i)} + \dots + \lambda_{s_i}^{(i)}$ is a partition, such that for all i and j ,

$$\lambda_j^{(i)} \geq \lambda_{j+c_{i+1}}^{(i+1)} \quad \text{and} \quad \lambda_j^{(k)} \geq \lambda_{j+c_1}^{(1)}.$$

We define the size $|\Lambda|$ of a cylindric partition $\Lambda = (\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$ to be the sum of all the parts in the partitions $\lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)}$. We also define the largest part of a cylindric partition Λ to be the maximum

part among all the partitions in Λ , and denote it by $\max(\Lambda)$. We now define the bivariate generating function for cylindric partitions

$$F_c(z, q) := \sum_{\Lambda \in \mathcal{P}_c} z^{\max(\Lambda)} q^{|\Lambda|},$$

where \mathcal{P}_c denotes the set of all cylindric partitions with profile c .

In 2007, Borodin [Bor07] showed that when one sets $z = 1$ in this bivariate generating function, it becomes a beautiful infinite product.

Theorem 2.1 (Borodin, 2007). *Let k and ℓ be positive integers, and let $c = (c_1, c_2, \dots, c_k)$ be a composition of ℓ . Define $t := k + \ell$ and $s(i, j) := c_i + c_{i+1} + \dots + c_j$. Then,*

$$F_c(1, q) = \frac{1}{(q^t; q^t)_\infty} \prod_{i=1}^k \prod_{j=i}^k \prod_{m=1}^{c_i} \frac{1}{(q^{m+j-i+s(i+1,j)}; q^t)_\infty} \prod_{i=2}^k \prod_{j=2}^i \prod_{m=1}^{c_i} \frac{1}{(q^{t-m+j-i-s(j,i-1)}; q^t)_\infty}.$$

In other words, if we denote by μ the partition $s(1, 1) + s(1, 2) + \dots + s(1, k)$, and by μ^c its complement inside the $k \times s(1, k)$ rectangle, we have

$$F_c(1, q) = \frac{1}{(q^t; q^t)_\infty} \prod_{\square \in \mu} \frac{1}{(q^{h(\square)}; q^t)_\infty} \prod_{\square \in \mu^c} \frac{1}{(q^{t-h(\square)}; q^t)_\infty},$$

where $h(\square)$ denotes the hook length of the box \square , see Figure 1.




FIGURE 1. The Borodin formula in terms of hooks

Theorem 2.1 is very useful for computing generating functions for cylindric partitions. For example, using the hooks given in Figure 2, we can easily obtain that the generating function for cylindric partitions with profile $(3, 1, 1)$ is

$$F_{(3,1,1)}(1, q) = \frac{1}{(q; q)_\infty} \times \frac{1}{(q, q, q^3, q^3, q^5, q^5, q^7, q^7; q^8)_\infty}.$$




FIGURE 2. The Borodin formula for profile $(3, 1, 1)$

Note that using the definition of cylindric partitions it is trivial to check that

$$F_c(z, q) = F_{S(c)}(z, q)$$

where $S(c) = (c_2, \dots, c_k, c_1)$. In the case $k = 3$, we have an extra identity.

Corollary 2.2. *For any positive integers c_1, c_2, c_3 , we have*

$$F_{(c_1, c_2, c_3)}(1, q) = F_{(c_2, c_1, c_3)}(1, q).$$




FIGURE 3. The Borodin formula for profiles (c_1, c_2, c_3) and (c_2, c_1, c_3)

Proof: We can assume without loss of generality that $c_1 > c_2$. In Figure 3, we can see that the hooks in the plain colored areas are exactly the same for both profiles.

It remains to see what happens in the dashed areas. First, we have

$$\prod_{\square \in C} \frac{1}{(q^{h(\square)}; q^t)_\infty} = \prod_{\square \in E} \frac{1}{(q^{t-h(\square)}; q^t)_\infty}.$$

Indeed, $\{h(\square) | \square \in C\} = \{c_2 + c_3 + 3, \dots, c_1 + c_3 + 2\}$. On the other hand, $\{h(\square) | \square \in E\} = \{c_2 + 1, \dots, c_1\}$, so that $\{t - h(\square) | \square \in E\} = \{c_2 + c_3 + 3, \dots, c_1 + c_3 + 2\}$.

Finally, the only thing left to do is proving that

$$(2.1) \quad \prod_{\square \in B} \frac{1}{(q^{h(\square)}; q^t)_\infty} \prod_{\square \in A} \frac{1}{(q^{t-h(\square)}; q^t)_\infty} = \prod_{\square \in F} \frac{1}{(q^{h(\square)}; q^t)_\infty} \prod_{\square \in D} \frac{1}{(q^{t-h(\square)}; q^t)_\infty}.$$

Indeed, we have:

$$\begin{aligned} \{h(\square) | \square \in B\} &= \{c_2 + 2, \dots, c_1 + 1\}, \\ \{t - h(\square) | \square \in A\} &= \{t - (c_2 + 2), \dots, t - (c_2 + c_3 + 1)\} \\ &= \{c_1 + 2, \dots, c_1 + c_3 + 1\}. \end{aligned}$$

On the side of the (c_2, c_1, c_3) profile, we have:

$$\begin{aligned} \{h(\square) | \square \in F\} &= \{c_2 + c_3 + 2, \dots, c_1 + c_3 + 1\}, \\ \{t - h(\square) | \square \in D\} &= \{t - (c_1 + 2), \dots, t - (c_1 + c_3 + 1)\} \\ &= \{c_2 + 2, \dots, c_2 + c_3 + 1\}. \end{aligned}$$

Thus $\{h(\square) | \square \in B\} \cup \{t - h(\square) | \square \in A\} = \{h(\square) | \square \in F\} \cup \{t - h(\square) | \square \in D\}$ and (2.1) is proved. \square

Corteel and Welsh [CW19] showed that $F_c(z, q)$ satisfies a nice q -difference equation. Let $c = (c_1, \dots, c_k)$ and use the convention that $c_0 = c_k$. Denote by I_c the set of indices $j \in \{1, \dots, k\}$ such that $c_j > 0$. Given a subset J of I_c , the composition $c(J) = (c_1(J), \dots, c_k(J))$ is defined by:

$$c_i(J) := \begin{cases} c_i - 1 & \text{if } i \in J \text{ and } i - 1 \notin J, \\ c_i + 1 & \text{if } i \notin J \text{ and } i - 1 \in J, \\ c_i & \text{otherwise.} \end{cases}$$

The q -difference equation is as follows.

Theorem 2.3 (Corteel–Welsh). *For any profile c ,*

$$F_c(z, q) = \sum_{\emptyset \subset J \subseteq I_c} (-1)^{|J|-1} \frac{F_{c(J)}(zq^{|J|}, q)}{(1 - zq^{|J|})},$$

with the initial conditions $F_c(0, q) = F_c(z, 0) = 1$.

Writing, for every composition c ,

$$G_c(z, q) := (zq; q)_\infty F_c(z, q),$$

Theorem 2.3 can be rewritten in the following form given in [CW19].

Theorem 2.4 (Cortee–Welsh, 2019). *For any profile c ,*

$$G_c(z, q) = \sum_{\emptyset \subset J \subseteq I_c} (-1)^{|J|-1} (zq; q)_{|J|-1} G_{c(J)}(zq^{|J|}, q),$$

with the initial conditions $G_c(0, q) = G_c(z, 0) = 1$.

Cylindric partitions, and in particular Theorems 2.1 and 2.4, are a very interesting tool to discover and prove Rogers–Ramanujan type identities.

Before we prove our new results, we present a warm-up exercise that gives a quick proof of the Rogers–Ramanujan identities, as in [Cor17]. These identities can be proved quite easily by using cylindric partitions with profiles $(3, 0)$ and $(2, 1)$. By Theorem 2.1, we have

$$(2.2) \quad G_{(3,0)}(1, q) = \frac{1}{(q^2, q^3; q^5)_\infty},$$

$$(2.3) \quad G_{(2,1)}(1, q) = \frac{1}{(q, q^4; q^5)_\infty}.$$

By Theorem 2.4, we also have

$$(2.4) \quad G_{(3,0)}(z, q) = G_{(2,1)}(zq, q),$$

$$(2.5) \quad G_{(2,1)}(z, q) = G_{(3,0)}(zq, q) + G_{(2,1)}(zq, q) - (1 - zq)G_{(2,1)}(zq^2, q).$$

Substituting (2.4) into (2.5), we obtain

$$G_{(2,1)}(z, q) = G_{(2,1)}(zq, q) + zqG_{(2,1)}(zq^2, q).$$

This equation is a classical linear second-order q -difference equation, and can be found for example in [And84b, eq. (7.1.1)]. Then we can easily show that

$$G_{(2,1)}(z, q) = \sum_{n \geq 0} \frac{z^n q^{n^2}}{(q; q)_n},$$

and deduce from (2.4) that

$$G_{(3,0)}(z, q) = \sum_{n \geq 0} \frac{z^n q^{n^2+n}}{(q; q)_n}.$$

Combining this with (2.2) and (2.3) proves the Rogers–Ramanujan identities.

This method was crucial in the recent paper of Cortee and Welsh [CW19], where they studied all cylindric partitions profiles where $k = 3$ and $\ell = 4$ to reprove the A_2 Rogers–Ramanujan identities due to Andrews, Schilling and Warnaar [ASW99].

In this paper, we use Theorems 2.1 and 2.4 on cylindric partitions with all profiles (c_1, c_2, c_3) such that $c_1 + c_2 + c_3 = 5$, to prove new A_2 Rogers–Ramanujan type identities. Our results can be stated as follows.

Theorem 2.5. *We have*

$$\begin{aligned} G_{(5,0,0)}(1, q) &= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2+n_2^2+n_3^2+n_4^2+n_1+n_2+n_3+n_4-n_1n_2+n_2n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\ &= \frac{1}{(q^2, q^3, q^3, q^4, q^4, q^5, q^5, q^6; q^8)_\infty}, \end{aligned}$$

$$\begin{aligned} G_{(4,1,0)}(1, q) &= G_{(4,0,1)}(1, q) \\ &= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2+n_2^2+n_3^2+n_4^2+n_2+n_3+n_4-n_1n_2+n_2n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \end{aligned}$$

$$\begin{aligned}
&= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_3 - n_1 n_2 + n_2 n_4} (1 + q^{n_1 + n_2 + n_4 + 1})}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\
&= \frac{1}{(q, q^2, q^3, q^4, q^4, q^5, q^6, q^7; q^8)_\infty},
\end{aligned}$$

$$\begin{aligned}
G_{(3,0,2)}(1, q) &= G_{(3,2,0)}(1, q) \\
&= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 - n_1 n_2 + n_2 n_4} (1 + q^{n_1 + n_3 + 1} + q^{2n_1 + n_2 + n_3 + n_4 + 2})}{(q; q)_{n_1}} \\
&\quad \times \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\
&= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 - n_1 n_2 + n_2 n_4} (q^{n_3} + q^{n_1 + 1} + q^{2n_1 + n_3 + 2} + q^{3n_1 + n_2 + n_3 + n_4 + 3})}{(q; q)_{n_1}} \\
&\quad \times \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\
&= \frac{1}{(q, q^2, q^2, q^3, q^5, q^6, q^6, q^7; q^8)_\infty},
\end{aligned}$$

$$\begin{aligned}
G_{(3,1,1)}(1, q) &= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_3 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\
&= \frac{1}{(q, q, q^3, q^3, q^5, q^5, q^7, q^7; q^8)_\infty},
\end{aligned}$$

$$\begin{aligned}
G_{(2,2,1)}(1, q) &= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\
&= \frac{1}{(q, q, q^2, q^4, q^4, q^6, q^7, q^7; q^8)_\infty}.
\end{aligned}$$

3. PROOF OF THEOREM 2.5

In this section, we prove our Rogers–Ramanujan type identities. For the product side, a direct application of Theorem 2.1 gives the following.

Corollary 3.1. *We have*

$$\begin{aligned}
G_{(5,0,0)}(1, q) &= \frac{1}{(q^2, q^3, q^3, q^4, q^4, q^5, q^5, q^6; q^8)_\infty}, \\
G_{(4,1,0)}(1, q) &= G_{(4,0,1)}(1, q) = \frac{1}{(q, q^2, q^3, q^4, q^4, q^5, q^6, q^7; q^8)_\infty}, \\
G_{(3,0,2)}(1, q) &= G_{(3,2,0)}(1, q) = \frac{1}{(q, q^2, q^2, q^3, q^5, q^6, q^6, q^7; q^8)_\infty}, \\
G_{(3,1,1)}(1, q) &= \frac{1}{(q, q, q^3, q^3, q^5, q^5, q^7, q^7; q^8)_\infty}, \\
G_{(2,2,1)}(1, q) &= \frac{1}{(q, q, q^2, q^4, q^4, q^6, q^7, q^7; q^8)_\infty}.
\end{aligned}$$

We will now use Theorem 2.4 to prove the sum sides. Our main result is the following expressions for $G_c(z, q)$ as quadruple series.

Theorem 3.2. *We have*

(3.1)

$$G_{(5,0,0)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q,$$

(3.2)

$$G_{(4,1,0)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q,$$

(3.3)

$$G_{(4,0,1)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_3 - n_1 n_2 + n_2 n_4} (1 + zq^{n_1 + n_2 + n_4 + 1})}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q$$

(3.4)

$$G_{(3,0,2)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 - n_1 n_2 + n_2 n_4} (1 + zq^{n_1 + n_3 + 1} + zq^{2n_1 + n_2 + n_3 + n_4 + 2})}{(q; q)_{n_1}} \times \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q,$$

(3.5)

$$G_{(3,2,0)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 - n_1 n_2 + n_2 n_4} (q^{n_3} + zq^{n_1 + 1} + zq^{2n_1 + n_3 + 2} + zq^{3n_1 + n_2 + n_3 + n_4 + 3})}{(q; q)_{n_1}} \times \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q,$$

(3.6)

$$G_{(3,1,1)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_3 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q,$$

(3.7)

$$G_{(2,2,1)}(z, q) = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{z^{n_1} q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q.$$

Note that even though, when $z = 1$, $G_{(4,1,0)}(1, q) = G_{(4,0,1)}(1, q)$ and $G_{(3,0,2)}(1, q) = G_{(3,2,0)}(1, q)$, in general $G_{(4,1,0)}(z, q)$ is not equal to $G_{(4,0,1)}(z, q)$ and $G_{(3,0,2)}(z, q)$ is not equal to $G_{(3,2,0)}(z, q)$.

Proof: Applying Theorem 2.4, we obtain the following q -difference equations for the compositions under consideration.

$$(3.8) \quad G_{(5,0,0)}(z, q) = G_{(4,1,0)}(zq, q),$$

$$(3.9) \quad G_{(4,1,0)}(z, q) = G_{(4,0,1)}(zq, q) + G_{(3,2,0)}(zq, q) - (1 - zq)G_{(3,1,1)}(zq^2, q),$$

$$(3.10) \quad G_{(4,0,1)}(z, q) = G_{(5,0,0)}(zq, q) + G_{(3,1,1)}(zq, q) - (1 - zq)G_{(4,1,0)}(zq^2, q),$$

$$(3.11) \quad G_{(3,2,0)}(z, q) = G_{(3,1,1)}(zq, q) + G_{(3,0,2)}(zq, q) - (1 - zq)G_{(2,2,1)}(zq^2, q),$$

$$(3.12) \quad G_{(3,1,1)}(z, q) = G_{(4,1,0)}(zq, q) + G_{(3,0,2)}(zq, q) + G_{(2,2,1)}(zq, q) \\ - (1 - zq)(G_{(4,0,1)}(zq^2, q) + G_{(3,2,0)}(zq^2, q) + G_{(2,2,1)}(zq^2, q)) \\ + (1 - zq)(1 - zq^2)G_{(3,1,1)}(zq^3, q),$$

$$(3.13) \quad G_{(3,0,2)}(z, q) = G_{(4,0,1)}(zq, q) + G_{(2,2,1)}(zq, q) - (1 - zq)G_{(3,1,1)}(zq^2, q),$$

$$(3.14) \quad G_{(2,2,1)}(z, q) = G_{(3,2,0)}(zq, q) + G_{(3,1,1)}(zq, q) + G_{(2,2,1)}(zq, q) \\ - (1 - zq)(G_{(3,1,1)}(zq^2, q) + G_{(3,0,2)}(zq^2, q) + G_{(2,2,1)}(zq^2, q)) \\ + (1 - zq)(1 - zq^2)G_{(2,2,1)}(zq^3, q).$$

We use Ablinger and Uncu's `qFunctions` package [AU19] together with Koutschan's `HolonomicFunctions` package [Kou09] to prove (3.1), (3.2), (3.6) and (3.7). The coupled q -difference equation system (3.8)-(3.14) corresponds to a coupled q -recurrence system. To that end, for any profile c we define the formal

$$G_c(z, q) = \sum_{k \geq 0} g_c(k) z^k,$$

where $g_c(k)$ is a function of q . The initial conditions $g_c(0) = 1$ and $g_c(k) = 0$ for every $k \leq 0$ define these sequences uniquely for all the relevant profiles c . Then, for example (3.9) is equivalent to

$$(3.15) \quad g_{(4,1,0)}(k) = q^k g_{(4,0,1)}(k) + q^k g_{(3,2,0)}(k) - q^{2k} g_{(3,1,1)}(k) + q^{2k-1} g_{(3,1,1)}(k-1),$$

for all $k \in \mathbb{Z}$. Uncoupling a coupled system of recurrences corresponds to Gaussian elimination and can be done automatically by symbolic computation implementations. Here we use the `HolonomicFunctions` package [Kou09], which uses Gröbner bases calculations to uncouple a given linear system of recurrences. Uncoupling the recurrences shows that the coefficients $g_{(5,0,0)}(k)$, $g_{(4,1,0)}(k)$, $g_{(3,2,0)}(k)$, $g_{(3,1,1)}(k)$, and $g_{(2,2,1)}(k)$ each satisfy recurrences of order 4 and $g_{(4,0,1)}(k)$ and $g_{(3,0,2)}(k)$ satisfy recurrences of order 6. We present the recurrence that $g_{(4,1,0)}(k)$ satisfies as an explicit example:

$$(3.16) \quad a_k(q)g_{(4,1,0)}(k) - b_k(q)g_{(4,1,0)}(k-1) - c_k(q)g_{(4,1,0)}(k-2) + d_k(q)g_{(4,1,0)}(k-3) + e_k(q)g_{(4,1,0)}(k-4) = 0,$$

where

$$\begin{aligned} a_k(q) &= (1 - q^k)(1 - q^{k-2} - 2q^{k-1} - q^k + q^{2k-3} + q^{2k-2} + q^{2k-1} - q^{3k-5} - q^{3k-4} - q^{3k-3} \\ &\quad - q^{4k-5} - q^{4k-4} - q^{4k-3} + q^{5k-7} + q^{5k-6} + q^{5k-5} - q^{6k-8} - q^{6k-7} - q^{6k-6}), \\ b_k(q) &= q^{2k+1}(1 + q^{k-2} - q^{2k-6} - 2q^{2k-5} - 4q^{2k-4} - 5q^{2k-3} - 5q^{2k-2} - 2q^{2k-1} - q^{2k} + q^{3k-7} + 2q^{3k-6} \\ &\quad + 4q^{3k-5} + 5q^{3k-4} + 6q^{3k-3} + 5q^{3k-2} + 3q^{3k-1} + q^{3k} - q^{4k-8} - 4q^{4k-7} - 6q^{4k-6} - 8q^{4k-5} \\ &\quad - 7q^{4k-4} - 7q^{4k-3} - 4q^{4k-2} - 2q^{4k-1} + q^{5k-8} + 2q^{5k-7} + 2q^{5k-6} + q^{5k-5} - q^{5k-2} + q^{6k-9} \\ &\quad + q^{6k-8} + 2q^{6k-7} + 2q^{6k-6} + 3q^{6k-5} + 3q^{6k-4} + 3q^{6k-3} + 2q^{6k-2} - q^{7k-10} - 3q^{7k-9} \\ &\quad - 5q^{7k-8} - 7q^{7k-7} - 7q^{7k-6} - 7q^{7k-5} - 5q^{7k-4} - 4q^{7k-3} - q^{7k-2} + q^{8k-10} + 3q^{8k-9} \\ &\quad + 5q^{8k-8} + 6q^{8k-7} + 4q^{8k-6} + 3q^{8k-5} + q^{8k-4} + q^{8k-3} - q^{9k-10} - 2q^{9k-9} - 3q^{9k-8} \\ &\quad - 2q^{9k-7} - q^{9k-6} - q^{10k-9} - q^{10k-8} - 2q^{10k-7} - q^{10k-6} - q^{10k-5} + q^{11k-10} + 2q^{11k-9} \\ &\quad + 2q^{11k-8} + q^{11k-7} - q^{12k-10} - q^{12k-9} - q^{12k-8}), \\ c_k(q) &= q^{3k-2}(1 + q - 2q^{k-1} - 3q^k - 2q^{k+1} - q^{k+2} + q^{2k-2} + 2q^{2k-1} + 2q^{2k} + q^{2k+1} + q^{2k+2} - q^{3k-5} - q^{3k-4} \\ &\quad - 3q^{3k-3} - 3q^{3k-2} - 3q^{3k-1} - q^{3k} - q^{3k+1} + q^{4k-7} + 2q^{4k-6} + 3q^{4k-5} + 3q^{4k-4} + 3q^{4k-3} \\ &\quad + q^{4k-2} - 2q^{4k} - q^{4k+1} - q^{4k+2} - q^{5k-8} - 2q^{5k-7} - 3q^{5k-6} - 2q^{5k-5} - 3q^{5k-4} - q^{5k-3} \\ &\quad + 2q^{5k-1} + 2q^{5k} + q^{5k+1} + q^{6k-9} + 3q^{6k-8} + 4q^{6k-7} + 4q^{6k-6} + 2q^{6k-5} + q^{6k-4} - 2q^{6k-3} \\ &\quad - 2q^{6k-2} - 4q^{6k-1} - 2q^{6k} - q^{6k+1} - q^{7k-9} - q^{7k-8} - q^{7k-7} + q^{7k-6} + 2q^{7k-5} + 3q^{7k-4} + 3q^{7k-3} \\ &\quad + 2q^{7k-2} + q^{7k-1} + q^{8k-9} - 2q^{8k-6} - 2q^{8k-5} - 3q^{8k-4} - 2q^{8k-3} - q^{8k-2} + q^{9k-9} + 2q^{9k-8} \\ &\quad + 3q^{9k-7} + 3q^{9k-6} + 2q^{9k-5} + q^{9k-4} - q^{10k-9} - q^{10k-8} - q^{10k-7} + q^{11k-9} + q^{11k-8} + q^{11k-7}), \\ d_k(q) &= q^{5k-4}(1 - q^{k-3} - q^{k-2} - q^{k-1} + q^{2k-7} + q^{2k-3} - q^{2k} - q^{3k-8} - 2q^{3k-7} - 2q^{3k-6} - 2q^{3k-5} - 2q^{3k-4} \\ &\quad - q^{3k-3} + q^{3k-1} + q^{3k} + q^{4k-8} + q^{4k-7} + q^{4k-6} - q^{4k-3} - q^{4k-2} - q^{4k-1} - q^{5k-9} - q^{5k-8} \\ &\quad - q^{5k-7} - q^{5k-6} - q^{5k-5} - q^{5k-4} - q^{6k-8} - q^{6k-7} - q^{6k-6} - q^{6k-5} - q^{6k-4} - q^{6k-3} + q^{7k-9} \\ &\quad + q^{7k-8} + q^{7k-7} - q^{8k-9} - q^{8k-8} - q^{8k-7} - q^{8k-6} - q^{8k-5} - q^{8k-4}), \\ e_k(q) &= q^{6k-9}(1 - q^{k-1} - 2q^k - q^{k+1} + q^{2k-1} + q^{2k} + q^{2k+1} - q^{3k-2} - q^{3k-1} - q^{3k} - q^{4k-1} - q^{4k} - q^{4k+1} \\ &\quad + q^{5k-2} + q^{5k-1} + q^{5k} - q^{6k-2} - q^{6k-1} - q^{6k}). \end{aligned}$$

The recurrence (3.16) alone shows that symbolic computation implementations such as `HolonomicFunctions` [Kou09] is a necessity for modern applications to perform tedious calculations with ease and eliminate any errors. Generating the coupled system of q -difference equations (3.8)-(3.14), turning these q -difference equations into q -recurrences in the form of (3.15) and uncoupling of the recurrence system (implicitly using `HolonomicFunctions`) can be automatically done by the `qFunctions` [AU19] package.

Now we can turn our attention on the claimed explicit expression (3.2) for $G_{(4,1,0)}(z, q)$. We can find a recurrence that the coefficients

$$g_k(q) := \sum_{n_2, n_3, n_4 \geq 0} \frac{q^{k^2 + n_2^2 + n_3^2 + n_4^2 + n_2 + n_3 + n_4 - kn_2 + n_2 n_4}}{(q; q)_k} \begin{bmatrix} k \\ n_2 \end{bmatrix}_q \begin{bmatrix} k \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q$$

of z^k in (3.2) satisfy using `HolonomicFunctions` [Kou09]. This is done by a successive application of Zeilberger's creative telescoping algorithm. Here one sees that $g_k(q)$ satisfies the same recurrence (3.16) that $g_{(4,1,0)}(k)$ satisfies. We can easily check that $g_k(q)$ and $g_{(4,1,0)}(k)$ satisfy the same initial conditions. This proves (3.2).

Same steps can be taken to prove (3.1), (3.6), and (3.7). In these cases, the uncoupled recurrences that come from (3.8)-(3.14) and the recurrences we get from the right-hand sides of (3.1), (3.6), and (3.7) coincide. Checking the trivial initial conditions proves these claims.

In the remaining cases, (3.3), (3.4), and (3.5), we cannot directly apply the available symbolic computation implementations. This is due to the extra polynomial factors that appear in these expressions. Technically speaking, we can still split these sums and find recurrences for each piece. Later we can use the closure properties of holonomic sequences under addition to find a recurrence that is satisfied by all pieces simultaneously. This gives us a recurrence of much higher order than necessary. These recurrences do not match the recurrences we find from the coupled system (3.8)-(3.14). Therefore, to finish the proofs this way, we would need to use the closure properties once again. This is complicated and it can be avoided all together.

With the identities (3.2), (3.6), and (3.7) proven, we now turn to the remaining sum formulas of Theorem 3.2. The formula (3.1) follows easily from (3.2) and (3.8).

Replacing z by zq in (3.8), we have

$$G_{(5,0,0)}(zq, q) = G_{(4,1,0)}(zq^2, q).$$

Substituting this into (3.10) yields

$$(3.17) \quad G_{(4,0,1)}(z, q) = G_{(3,1,1)}(zq, q) + zqG_{(4,1,0)}(zq^2, q).$$

Using (3.2) and (3.6) in this new equation (3.17), we prove (3.3).

Now in (3.13), we replace $G_{(4,0,1)}(zq, q)$ by its expression given by (3.17), and obtain

$$(3.18) \quad G_{(3,0,2)}(z, q) = G_{(2,2,1)}(zq, q) + zqG_{(3,1,1)}(zq^2, q) + zq^2G_{(4,1,0)}(zq^3, q).$$

So we can prove (3.4) by using (3.2), (3.6), and (3.7) in (3.18).

Finally, in (3.11), we replace $G_{(3,0,2)}(zq, q)$ by its expression given by (3.18). This gives

$$(3.19) \quad G_{(3,2,0)}(z, q) = G_{(3,1,1)}(zq, q) + zqG_{(2,2,1)}(zq^2, q) + zq^2G_{(3,1,1)}(zq^3, q) + zq^3G_{(4,1,0)}(zq^4, q).$$

Thus we can prove the final sum (3.5) by using (3.2), (3.6), and (3.7) in (3.19). \square

Combining Theorems 3.1 and 3.2 concludes the proof of Theorem 2.5.

4. CONCLUSION

In this note we made a first progress towards the general family of A_2 Rogers–Ramanujan identities. To get the full picture, we would need to compute the generating functions of cylindric partitions of profile (c_1, c_2, c_3) for any $c_1, c_2, c_3 \in \mathbb{N}$. The Rogers–Ramanujan identities correspond to the case $c_1 + c_2 + c_3 = 2$. The previously known A_2 Rogers–Ramanujan identities of Andrews, Schilling, and Warnaar correspond to the case $c_1 + c_2 + c_3 = 4$. In this note we settle the case $c_1 + c_2 + c_3 = 5$. For now we do not have the right tools to conjecture the sum side for general (c_1, c_2, c_3) and we do not know how to go further. Nevertheless

the problem seems to have some beautiful structure. Let us define $f_{n,c}(q)$ to be the generating function of cylindric partitions with profile c with entries bounded by n . Or equivalently,

$$f_{n,c}(q) := [z^n] \frac{F_c(z, q)}{1-z}.$$

And let $g_{n,c} := [z^n] G_c(z, q)$. For example, using the combinatorics of lattice paths, one can prove the following.

Lemma 4.1. *The generating function of cylindric partitions with profile $(\ell, 0, 0)$ with entries bounded by n is*

$$f_{n,(\ell,0,0)}(q) = \frac{P_{n,(\ell,0,0)}(q)}{(q^3; q^3)_n},$$

where P is a polynomial in q with integer coefficients and $P_{n,(\ell,0,0)}(1) = \binom{\ell+2}{2}^n$.

See [GK97] to see how to write cylindric partitions as non-intersecting paths.

Using Theorem 2.3, we can prove that this holds in more generality. The generating function for cylindric partitions with profile $c = (c_1, c_2, c_3)$ with entries bounded by n is

$$f_{n,c}(q) = \frac{P_{n,c}(q)}{(q^3; q^3)_n},$$

where P is a polynomial in q with integer coefficients and $P_{n,c}(1) = \binom{\ell+2}{2}^n$, where $\ell = c_1 + c_2 + c_3$. We conjecture that

Conjecture 4.2. *For any profile $c = (c_1, c_2, c_3)$, the polynomial $P_{n,c}(q)$ has non-negative coefficients. Moreover if $c_1 + c_2 + c_3 \not\equiv 0 \pmod{3}$, the generating function $g_{n,c}$ can be written as*

$$g_{n,c} = \frac{1}{(q; q)_n} Q_{n,c}(q),$$

where $Q_{n,c}(q)$ is a polynomial in q with non-negative coefficients and $Q_{n,c}(1) = \left(\frac{(\ell+2)(\ell+1)}{6} - 1 \right)^n$, where $\ell = c_1 + c_2 + c_3$.

This conjecture is true for any n if $c_1 + c_2 + c_3 = 2$ (Rogers–Ramanujan identities), $c_1 + c_2 + c_3 = 4$ (Andrews–Schilling–Warnaar A_2 Rogers–Ramanujan identities) and $c_1 + c_2 + c_3 = 5$ (this paper). For example

$$Q_{n,(2,2,1)}(q) = \sum_{n_2, n_3, n_4 \geq 0} q^{n^2 + n_2^2 + n_3^2 + n_4^2 - n n_2 + n_2 n_4} \begin{bmatrix} n \\ n_2 \end{bmatrix}_q \begin{bmatrix} n \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q$$

and $Q_{n,(2,2,1)}(1) = 6^n$. This conjecture has been checked for small values of k and n .

We know by Corollary 2.2 and by definition of cylindric partitions that $G_{(c_1, c_2, c_3)}(1, q)$ is the same for any permutation of the c_i 's. From this, we directly deduce that $G_{(4,0,1)}(1, q) = G_{(4,1,0)}(1, q)$, i.e.

$$\begin{aligned} \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q \\ = \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_3 - n_1 n_2 + n_2 n_4} (1 + q^{n_1 + n_2 + n_4 + 1})}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q. \end{aligned}$$

The referee suggested that it is also possible to obtain this q -series identity directly by showing that

$$\sum_{n_1, n_4 \geq 0} \frac{q^{n_1^2 + n_4^2 - n_1 n_2 + n_2 n_4 + n_2 + n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q = \sum_{n_1, n_4 \geq 0} \frac{q^{n_1^2 + n_4^2 - n_1 n_2 + n_2 n_4 + n_1}}{(q; q)_{n_1}} (1 + q^{n_1 + n_2 + n_4 + 1}) \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q$$

holds for all non-negative integer n_2 .

Finally, we mentioned in the Introduction that Theorem 5.3 of Andrews, Schilling, and Warnaar [ASW99] gives another expression of all the sums in our Theorem 2.5, but their sums have a factor $(q; q)_\infty$ in front of them, like in Theorem 1.5. So far, the only way we know that their sums and ours are equal is also because

they equal the same infinite products. It would be an interesting exercise to find a direct connection between the sums as well. For example, can one find a q -series proof of the equality

$$(q, q)_\infty \sum_{a_1, b_1, a_2, b_2 \in \mathbb{Z}} \frac{q^{a_1^2 + b_1^2 + a_2^2 + b_2^2 - a_1 b_1 + a_2 b_2 + a_1 + a_2 + b_1 + b_2}}{(q; q)_{a_1 - a_2} (q; q)_{b_1 - b_2} (q; q)_{a_2} (q; q)_{b_2} (q; q)_{a_2 + b_2 + 1}}$$

$$= \sum_{n_1, n_2, n_3, n_4 \geq 0} \frac{q^{n_1^2 + n_2^2 + n_3^2 + n_4^2 + n_1 + n_2 + n_3 + n_4 - n_1 n_2 + n_2 n_4}}{(q; q)_{n_1}} \begin{bmatrix} n_1 \\ n_2 \end{bmatrix}_q \begin{bmatrix} n_1 \\ n_4 \end{bmatrix}_q \begin{bmatrix} n_2 \\ n_3 \end{bmatrix}_q ?$$

REFERENCES

- [AAB87] A. Agrawal, G. E. Andrews, and D. Bressoud. The Bailey lattice. *J. Indian Math. Soc.*, 51:57–73, 1987.
- [And74] G. E. Andrews. An analytic generalization of the Rogers-Ramanujan identities for odd moduli. *Proc. Nat. Acad. Sci. USA*, 71:4082–4085, 1974.
- [And84a] G. E. Andrews. Multiple series Rogers-Ramanujan type identities. *Pacific J. Math*, 114:267–283, 1984.
- [And84b] G. E. Andrews. *The Theory of Partitions*. Cambridge University Press, 1984.
- [And89] G. E. Andrews. On the proofs of the Rogers-Ramanujan identities. In *q-Series and Partitions*, pages 1–14. Springer-Verlag, New York, 1989.
- [ASW99] G. E. Andrews, A. Schilling, and S. O. Warnaar. An A_2 Bailey lemma and Rogers-Ramanujan-type identities. *J. Amer. Math. Soc.*, 12(3):677–702, 1999.
- [AU19] J. Ablinger and A. K. Uncu. **qFunctions** - a Mathematica package for q -series and partition theory applications. Submitted. arXiv:1910.12410, 2019.
- [Bai49] W. N. Bailey. Identities of the Rogers-Ramanujan type. *Proc. London Math. Soc.*, 50(2):1–10, 1949.
- [Bax81] R. J. Baxter. Rogers-Ramanujan identities in the hard hexagon model. *J. Stat. Phys.*, 26:427–452, 1981.
- [BMS13] C. Bruschek, H. Mourtada, and J. Schepers. Arc spaces and Rogers-Ramanujan identities. *Ramanujan J.*, 30:9–38, 2013.
- [Bor07] A. Borodin. Periodic Schur process and cylindric partitions. *Duke Math. J.*, 140(3):391–468, 2007.
- [Bre79] D. M. Bressoud. A generalization of the Rogers-Ramanujan identities for all moduli. *J. Comb. Th. A*, 27:64–68, 1979.
- [Bre83] D. M. Bressoud. An easy proof of the Rogers-Ramanujan identities. *J. Number Th.*, 16:335–241, 1983.
- [Cor17] S. Corteel. Rogers-Ramanujan identities and the Robinson-Schensted-Knuth correspondence. *Proc. Amer. Math. Soc.*, 145(5):2011–2022, 2017.
- [CW19] S. Corteel and T. A. Welsh. The A_2 Rogers–Ramanujan identities revisited. *Annals of Combinatorics*, 23(3):683–694, 2019.
- [FFW08] B. Feigin, O. Foda, and T. A. Welsh. Andrews–Gordon type identities from combinations of Virasoro characters. *Ramanujan J.*, 17(1):33–52, 2008.
- [FW16] O. Foda and T. A. Welsh. Cylindric partitions, \mathcal{W}_r characters and the Andrews–Gordon–Bressoud identities. *J. Phys. A*, 49(16):164004, 37, 2016.
- [GK97] I. M. Gessel and C. Krattenthaler. Cylindric partitions. *Trans. Amer. Math. Soc.*, 349(2):429–479, 1997.
- [GM81] A. M. Garsia and S. C. Milne. A Rogers-Ramanujan bijection. *J. Combin. Theory Ser. A*, 31:289–339, 1981.
- [Gor61] B. Gordon. A combinatorial generalisation of the Rogers-Ramanujan identities. *Amer. J. Math.*, 83:393–399, 1961.
- [GOW16] M. J. Griffin, K. Ono, and S. O. Warnaar. A framework of Rogers–Ramanujan identities and their arithmetic properties. *Duke Math. J.*, 8:1475–1527, 2016.
- [Kou09] C. Koutschan. *Advanced applications of the holonomic systems approach*. PhD thesis, RISC, Johannes Kepler University, Linz, 2009.
- [LW84] J. Lepowsky and R. L. Wilson. The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities. *Invent. Math.*, 77:199–290, 1984.
- [LW85] J. Lepowsky and R. L. Wilson. The structure of standard modules, II: The case $A_1^{(1)}$, principal gradation. *Invent. Math.*, 79:417–442, 1985.
- [Mac16] P. A. MacMahon. *Combinatory Analysis*, volume 2. Cambridge University Press, New York, NY, USA, 1916.
- [ML92] S. C. Milne and G. M. Lilly. The A_ℓ and C_ℓ Bailey transform and lemma. *Bull. Amer. Math. Soc.*, 26:258–263, 1992.
- [ML95] S. C. Milne and G. M. Lilly. Consequences of the A_ℓ and C_ℓ Bailey transform and lemma. *Discrete Math.*, 139:319–346, 1995.
- [Pas20] A. Pascadi. Several new product identities in relation to two-variable Rogers–Ramanujan type sums and mock theta functions. arXiv:2009.05878, 2020.
- [RR19] L. J. Rogers and S. Ramanujan. Proof of certain identities in combinatory analysis. *Camb. Phil. Soc. Proc.*, 19:211–216, 1919.
- [Sch17] I. Schur. Ein Beitrag zur Additiven Zahlentheorie und zur Theorie der Kettenbrüche. *S.-B. Preuss. Akad. Wiss. Phys. Math. Klasse*, pages 302–321, 1917.
- [Sil17] A. V. Sills. *An invitation to the Rogers-Ramanujan identities*. CRC Press, 2017.
- [War06] S. O. Warnaar. Hall–Littlewood functions and the A_2 Rogers–Ramanujan identities. *Adv. Math.*, 200(2):403–434, 2006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, USA
Email address: `corteel@berkeley.edu`

UNIV LYON, CNRS, UNIVERSITÉ CLAUDE BERNARD LYON 1, UMR5208, INSTITUT CAMILLE JORDAN, F-69622 VILLEURBANNE,
FRANCE
Email address: `dousse@math.cnrs.fr`

JOHANN RADON INSTITUTE FOR COMPUTATIONAL AND APPLIED MATHEMATICS, AUSTRIAN ACADEMY OF SCIENCE, ALTENBERG-
ERSTRASSE 69, A-4040 LINZ, AUSTRIA
Email address: `akuncu@risc.jku.at`